



image

1635

TRANSMITTAL LETTER
(General - Patent Pending)

Docket No.
P23,461A USA

In Re Application Of: **June Kaplow, Thomas Haws, Marie Rosier and Patrice Deneffe**

Serial No.
09/823,119

Filing Date
March 30, 2001

Examiner
J. Epps

Group Art Unit
1635

Title: **NUCLEAR FACTOR KAPPA B INDUCING FACTOR**

TO THE COMMISSIONER FOR PATENTS:

Transmitted herewith is: **certified copy of U.K. priority Application No. 0018307.9, filed July 26, 2000**

in the above identified application.

- ☒ No additional fee is required.
- ☐ A check in the amount of _____ is attached.
- ☒ The Director is hereby authorized to charge and credit Deposit Account No. **19-5425** as described below.
- ☐ Charge the amount of _____
- ☒ Credit any overpayment.
- ☒ Charge any additional fee required.

Signature

Patrick J. Kelly
Synnestvedt & Lechner LLP
2600 Aramark Tower
1101 Market Street
Philadelphia, PA 19107-2950
Telephone: 215-923-4466
Facsimile: 215-923-2189
email: pkelly@synnlech.com

Dated: **September 17, 2003**

CC:

I certify that this document and fee is being deposited on **09/17/2003** with the U.S. Postal Service as first class mail under 37 C.F.R. 1.8 and is addressed to the Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

Signature of Person Mailing Correspondence

Katrina Lynn

Typed or Printed Name of Person Mailing Correspondence

17



INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed

Dated

9 September 2003

100

Patents Form 1/77

Patents Act 1977
(Rule 16)

26 JUL 2000

NEWPORT

The
Patent
Office

27JUL00 E555956-1 D02298
P01/7700 0:00-0018307.9

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road
Newport
Gwent NP9 1RH

1. Your reference A3539 US

2. Patent application number
(The Patent Office will fill in this part)

0018307.9

26 JUL 2000

3. Full name, address and postcode of the or of each applicant (underline all surnames)

AVENTIS PHARMACEUTICALS PRODUCTS INC
500 ARCOLA ROAD
COLLEGEVILLE
PENNSYLVANIA 19426

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

USA

7880883001

4. Title of the invention

POLYPEPTIDES

5. Name of your agent (if you have one)

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

Adams Jones
Broadway Business Centre
32a Storey Street
Nottingham
NG1 1LL

Patents ADP number (if you know it)

7939648001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number
(if you know it)

Date of filing
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

YES

- a) any applicant named in part 3 is not an inventor, or
 - b) there is an inventor who is not named as an applicant, or
 - c) any named applicant is a corporate body.
- See note (d))

Patents Form 1/77

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description	64
Claim(s)	32 5
Abstract	1
Drawing(s)	

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature

Date

ANDREW AGNEW, AGENT FOR THE APPLICANT

July 25, 2000

12. Name and daytime telephone number of person to contact in the United Kingdom

ANDREW AGNEW 01477 538508

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

POLYPEPTIDES

Field of the Invention

Nuclear factor B (NFB) comprises a family of eukaryotic transcription factors
5 involved in regulation of genes involved in immune responses and other cellular
functions. In some instances, NFB activation leads to an inflammatory response which
ultimately results in a disease state. Accordingly, it would be desirable to develop
means for controlling induction of NFB. The present invention provides polypeptides
which are involved in the induction of NFB and which may be used to enhance NFB
10 expression or activation or to identify and prepare inhibitors of NFB expression or
activation.

Reported Developments

Nuclear factor B (NFB) comprises a family of transcription factors found in
15 almost all eukaryotic cells. NFB plays a role in the regulation of genes involved in
tissue inflammation, cellular proliferation, and cellular differentiation.

A number of studies have examined the relationship between disease states and
expression of the subunit proteins which comprise NFB or activation of NFB already
present in a cell. Li et al. studied the regulation of NFB by the HTLV-1 Tax protein
20 demonstrating that the ability of Tax to activate the NFB pathway plays an essential role
in HTLV-1-induced cellular transformation (Li et al., *Gene Expr.*, 7, 4-6, 233-245
(1999)). Lentsch and Ward studied the activation and regulation of NFB during acute
inflammation and described the relationship between NFB and the inhibiting proteins of
the IKappaB family (Lentsch et al., *Clin. Chem. Lab. Med.*, 37, 3, 205-208 (1999)).
25 Visconti et al. investigated the role of NFB in thyroid carcinogenesis by analyzing
thyroid carcinoma cell lines. Their studies indicated that activation of the NFB complex
by overexpression of the p65 protein played a critical role in the process of thyroid cell
transformation (Visconti et al., *Oncogene*, 15, 16, 1987-1994 (1997)). Mukhopadhyay
et al. investigated the expression of the p50 subunit of NFB transcription factor complex
30 in non-small cell lung carcinoma tissues demonstrating that 81 % of fresh non-small cell
lung cancer tissues express from two to twenty-fold higher levels of the p50 subunit than

normal lung tissue (Mukhopadhyay et al., *Oncogene*, 11, 5, 999-1003 (1995)). Khaled et al. targeted p50 gene expression with specific antisense 3' phosphorothioate modified antisense oligodeoxynucleotides and were able to reduce NFB expression. Their results demonstrated p50 antisense molecules could reduce NFB expression and could down-regulate the immune response providing possible treatment for autoimmune disorders (Khaled et al., *Clin. Immunol. Immunopathol.*, 83, 3, 254-263 (1997)).

Summary of the Invention

In accordance with the present invention, there are provided nucleic acid sequences encoding nuclear factor B inducing factor (NFIF) 14b and 7a. Included also are cDNAs encoding NFIF-14b and NFIF-7a and isolated and purified NFIF-14b and NFIF-7a polypeptides which induce NFB. The present invention includes also methods of inducing NFB comprising introducing into the body of a patient a composition that activates NFB induction, including for example, expression vectors that express NFIF-14b and NFIF-7a polypeptides. Examples of preferred expression vectors are retroviral vectors, adenoviral vectors, adeno-associated viral vectors, herpes, herpesviral vectors and naked DNA vectors. The present invention also provides methods wherein the composition of the present invention comprise an NFIF-14b or NFIF-7a polypeptide and a pharmaceutically acceptable carrier.

Another aspect of the present invention are compositions for lowering the expression of the NFIF gene comprising an antisense nucleic acid. Still another aspect of the present invention is a composition for lowering the activity of an NFIF polypeptide comprising a neutralizing antibody that binds to an NFIF polypeptide and lowers its activity.

Yet another embodiment of the present invention is a composition for lowering the expression of NFIF in a patient comprising a ribozyme that cuts RNA encoding an NFIF polypeptide.

The present invention provides methods for evaluating whether a test compound is effective in inhibiting the activity of NFIF-14b polypeptides comprising (A)

comparing the level of NFB-regulated gene expression in a first sample comprising: (1) NFIF-14b; (2) the NFB-regulated reporter gene; and (3) the test compound with the level of gene expression in a second sample comprising (4) NFIF-14b; and (5) the NFB-regulated reporter gene; and (B) determining whether the expression of the reporter gene
5 is lower in the first sample relative to the second sample.

Still another aspect of the present invention provides methods for evaluating whether a test compound is effective in inhibiting the activity of NFIF-7a polypeptides comprising (A) comparing the level of NFB-regulated gene expression in a first sample
10 comprising: (1) NFIF-7a; (2) the NFB-regulated reporter gene; and (3) the test compound with the level of gene expression in a second sample comprising (4) NFIF-7a; and (5) the NFB-regulated reporter gene; and (B) determining whether the expression of the reporter gene is lower in the first sample relative to the second sample.

15 Yet another aspect of the present invention provides a method for identifying whether a test compound can enhance the activity of NFIF-14b based on the expression of an NFB regulator reporter gene comprising (A) comparing the level of NFB-regulated gene expression in a first sample comprising (1) NFIF-14b; (2) the NFB-regulated reporter gene; and (3) the test compound with the level of gene expression in a second
20 sample comprising: (4) NFIF-14b; and (5) the NFB-regulated reporter gene; and (B) determining whether the expression of the reporter gene is higher in the first sample relative to the second sample.

Another aspect of the present invention provides a method for identifying
25 whether a test compound can enhance the activity of NFIF-7a based on the expression of an NFB regulator reporter gene comprising (A) comparing the level of NFB-regulated gene expression in a first sample comprising (1) NFIF-7a; (2) the NFB-regulated reporter gene; and (3) the test compound with the level of gene expression in a second sample comprising: (4) NFIF-7a; and (5) the NFB-regulated reporter gene; and (B)
30 determining whether the expression of the reporter gene is higher in said first sample relative to the second sample.

In yet another aspect of the present invention, methods are provided for inhibiting expression of NFB-dependent genes comprising administering to a patient a composition that inhibits the activity of NFIF-14b or NFIF-7a.

5 Still yet another aspect of the present invention is a method of inhibiting inflammation comprising administration of a composition that inhibits the activity of NFIF-14b or NFIF-7a.

10 Another aspect of the present invention relates to the use of an NFIF polypeptide for the manufacture of a medicament intended for the treatment and/or prevention of an NFB-regulated inflammatory response. Still yet another aspect of the present invention relates to the use of a nucleic acid encoding an NFIF polypeptide, for the manufacture of a medicament intended for the treatment and/or prevention of an NFB-regulated inflammatory response.

15 Another aspect of the present invention relates to the use of a recombinant vector comprising a nucleic acid encoding an NFIF polypeptide, for the manufacture of a medicament intended for the treatment and/or prevention of an NFB-regulated inflammatory response. Yet another aspect of the present invention relates to the use of a defective recombinant viral vector for the manufacture of a medicament intended for the treatment and/or prevention of an NFB-regulated response.

20 Another aspect of the present invention relates to the use of cells genetically modified *ex vivo* with a recombinant virus or production of cells containing such recombinant viruses which are implanted in the body, facilitating prolonged and effective expression *in vivo* of an NFIF polypeptide according to the invention.

25 Two sequences demonstrating similarities to the polypeptides of the present invention are available in the GenBank nucleic acid sequence library. The first sequence, having Accession Number Y08135 encodes a peptide identified as full-length mouse acid sphingomyelinase-like phosphodiesterase 3a. The second sequence, having Accession Number Y08136 encoded a partial human clone (863 bp of which 536 were

coding) for a protein sequence identified as human acid sphingomyelinase-like phosphodiesterase 3a. Both sequences were submitted on September 17, 1996 by K. Hofmann. Although both of these sequences were identified as encoding acid sphingomyelinase-like phosphodiesterases, there has not been confirmation of this type of enzymatic activity. These sequences may have been identified as acid sphingomyelinase-like phosphodiesterases due to similarities between the 3' sequences of these proteins and the 3' sequences of members of the sphingomyelinase family.

Brief Description of the Drawings

10 Figure 1 is the amino acid sequence for NFIF-14b.

Figure 2 is the amino acid sequence for NFIF-7a.

Figure 3 is the cDNA sequence for NFIF-14b and NFIF-7a aligned for comparison.

Figure 4 is a diagrammatic representation of NFIF-14b and NFIF-7a.

15 Figure 5 is a bar graph representing luminometer counts per second observed in HeK 293 cells at 48 hours.

Figure 6 is a bar graph representing luminometer counts per second observed in Cos-7 cells at 48 hours.

Figure 7 is the amino acid sequence of a 15-residue sequence used to prepare antibodies directed against NFIF.

Figure 8 is a Northern Blot using an NFIF probe.

Detailed Description of the Invention

The present invention provides polypeptides which are capable of inducing NFB as well as compounds and compositions that are capable of inhibiting induction of NFB.

The present invention is based in part on the discovery of NFB-inducing factor (NFIF) proteins. Two functional variants of the NFIF protein have been identified. One variant, NFIF-14b, comprises full-length NFIF. The second variant, NFIF-7a, is believed to be a splice variant of NFIF-14b. Both NFIF-14b and NFIF-7a have the ability to induce NFB.

NFB comprises a family of eukaryotic transcription factors which regulate genes involved in immune responses and other cellular activities. In certain situations, it may be desirable to induce NFB in order to initiate or increase the extent of an immune response. In other situations, it may be desirable to reduce or prevent NFB induction in order to reduce or prevent an NFB-regulated immune response. For example, it has been discovered that NFIF is associated with a variety of pathologies associated with NFB-regulated immune responses, including atherosclerosis. By inhibiting the expression of the NFIF gene, or by otherwise interfering with the activity of the NFIF protein, it is possible to inhibit NFB induction, thereby inhibiting or preventing NFB-regulated inflammatory responses that result in atherosclerosis and other diseases.

Discovery of the gene encoding NFIF and the ability to prepare NFIF proteins facilitates identification and preparation of compounds and compositions which are capable of inhibiting the activity of NFIF proteins and consequently inhibiting NFB induction.

The description which follows begins with a Definitions section, followed by background information relating to NFB. This is followed by a discussion of the NFIF polypeptides of the present invention, as well as information on isolation of DNA encoding such polypeptides and methods for preparing variants of the polypeptides. Expression vectors capable of expressing the encoding NFIF polypeptide DNAs are then discussed, including information on appropriate promoters for the expression systems, methods for introducing the expression vectors into appropriate host cells and viral vector systems. Following this discussion of NFIF polypeptides and their variants, there is a discussion of therapeutic uses for NFIF polypeptides. This discussion is followed by a description of compositions useful in the practice of the present invention. This is followed by a discussion which outlines the various therapeutic compositions and compounds, including polypeptides based on NFIF, antisense nucleic acids, ribozymes and antibodies. Finally, the methods of the present invention which use the described compounds and compositions are described.

Definitions

The term NFIF@ is used to designate Nuclear factor B-inducing factor@.

The terms Ainducing@ or Ainduction@ when used in describing the activity of NFIF include within their scope the ability to cause expression of the subunit proteins which comprise NFB either directly or indirectly as well as the ability to activate NFB that is already present in a cell. This activation may be direct or indirect and results in NFB functioning as a transcription factor.

For the purposes of the present description, the expression Anucleotide sequence@ may be used to designate either a polynucleotide or a nucleic acid. The expression Anucleotide sequence@ covers the genetic material itself and is therefore not restricted to the information relating to its sequence.

The terms Anucleic acid@, Apolynucleotide@, Aoligonucleotide@ or Anucleotide sequence@ cover RNA, DNA, gDNA or cDNA sequences or alternatively RNA/DNA hybrid sequences of more than one nucleotide, either in the single-stranded form or in the duplex, double-stranded form.

A Anucleic acid@ is a polymeric compound comprised of covalently linked subunits called nucleotides. Nucleic acid includes polyribonucleic acid (RNA) and polydeoxyribonucleic acid (DNA), both of which may be single-stranded or double-stranded. DNA includes cDNA, genomic DNA, synthetic DNA, and semi-synthetic DNA. The sequence of nucleotides that encodes a protein is called the sense sequence or coding sequence.

25

The term Anucleotide@ designates both the natural nucleotides (A, T, G, C) as well as the modified nucleotides that comprise at least one modification such as (1) an analog of a purine, (2) an analog of a pyrimidine, or (3) an analogous sugar, examples of such modified nucleotides being described, for example, in the PCT application

30 No. WO 95/04064.

For the purposes of the present invention, a first polynucleotide is considered as being complementary to a second polynucleotide when each base of the first nucleotide is paired with the complementary base of the second polynucleotide whose orientation is reversed. The complementary bases are A and T (or A and U), or C and G.

"Heterologous" DNA refers to DNA not naturally located in the cell, or in a chromosomal site of the cell. Preferably, the heterologous DNA includes a gene foreign to the cell.

As used herein, the term "homologous" in all its grammatical forms and spelling variations refers to the relationship between proteins that possess a "common evolutionary origin," including proteins from superfamilies (*e.g.*, the immunoglobulin superfamily) and homologous proteins from different species (*e.g.*, myosin light chain, etc.) (Reeck et al., *Cell*, 50:667 (1987)). Such proteins (and their encoding genes) have sequence homology, as reflected by their high degree of sequence similarity.

Accordingly, the term "sequence similarity" in all its grammatical forms refers to the degree of identity or correspondence between nucleic acid or amino acid sequences of proteins that may or may not share a common evolutionary origin (*see* Reeck et al., *supra*). However, in common usage and in the instant application, the term "homologous," when modified with an adverb such as "highly," may refer to sequence similarity and not a common evolutionary origin.

In a specific embodiment, two DNA sequences are "substantially homologous" or "substantially similar" when at least about 50% (preferably at least about 75%, and more preferably at least about 90 or 95%) of the nucleotides match over the defined length of the DNA sequences. Sequences that are substantially homologous can be identified by comparing the sequences using standard software available in sequence data banks, or in a Southern hybridization experiment under, for example, stringent conditions as defined for that particular system. Defining appropriate hybridization conditions is within the skill of the art. See, *e.g.*, Maniatis et al., *Molecular cloning: A Laboratory Manual*,

Cold Spring Harbor Press, Cold Spring, NY (1982); Glover et al., *DNA Cloning: A Practical Approach, Volumes I and II Oligonucleotide Synthesis*, MRL Press, Ltd., Oxford, U.K. (1985); Hames and Higgins, Hames BD and Higgins SJ, 1985. *Nucleic acid hybridization: a practical approach*, Hames and Higgins Ed., IRL Press, Oxford (1985)).

Similarly, in a particular embodiment, two amino acid sequences are "substantially homologous" or "substantially similar" when greater than 30% of the amino acids are identical, or greater than about 60% are similar (functionally identical). Preferably, the similar or homologous sequences are identified by alignment using, for example, the GCG (Genetics Computer Group, Program Manual for the GCG Package, Version 7, Madison, Wisconsin) pileup program.

The Apercentage identity@ between two nucleotide or amino acid sequences, for the purposes of the present invention, may be determined by comparing two sequences aligned optimally, through a window for comparison.

The portion of the nucleotide or polypeptide sequence in the window for comparison may thus comprise additions or deletions (for example Agaps@) relative to the reference sequence (which does not comprise these additions or these deletions) so as to obtain an optimum alignment of the two sequences.

The percentage is calculated by determining the number of positions at which an identical nucleic base or an identical amino acid residue is observed for the two sequences (nucleic or peptide) compared, and then by dividing the number of positions at which there is identity between the two bases or amino acid residues by the total number of positions in the window for comparison, and then multiplying the result by 100 in order to obtain the percentage sequence identity.

The optimum sequence alignment for the comparison may be achieved using a computer with the aid of known algorithms contained in the package from the company

WISCONSIN GENETICS SOFTWARE PACKAGE, GENETICS COMPUTER
GROUP (GCG), 575 Science Doctor , Madison, WISCONSIN.

By way of illustration, it will be possible to produce the percentage sequence
5 identity with the aid of the BLAST software (versions BLAST 1.4.9 of March 1996,
BLAST 2.0.4 of February 1998 and BLAST 2.0.6 of September 1998), using
exclusively the default parameters (Altschul et al, *J. Mol. Biol.*, 215:403-410 (1990);
Altschul et al, *Nucleic Acids Res.*, 25:3389-3402 (1997)). Blast searches for sequences
similar/homologous to a reference Arequest@ sequence, with the aid of the Altschul et al.
10 algorithm. The request sequence and the databases used may be of the peptide or nucleic
types, any combination being possible.

The term "corresponding to" is used herein to refer to similar or homologous
sequences, whether the exact position is identical or different from the molecule to
15 which the similarity or homology is measured. A nucleic acid or amino acid sequence
alignment may include spaces. Thus, the term "corresponding to" refers to the sequence
similarity, and not the numbering of the amino acid residues or nucleotide bases.

AVariant@ of a nucleic acid according to the invention will be understood to
20 mean a nucleic acid that differs by one or more bases relative to the reference
polynucleotide. A variant nucleic acid may be of natural origin, such as an allelic variant
that exists naturally, or it may also be a non-natural variant obtained, for example, by
mutagenic techniques.

25 In general, the differences between the reference (generally, wild-type) nucleic
acid and the variant nucleic acid are small such that the nucleotide sequences of the
reference nucleic acid and of the variant nucleic acid are very similar and, in many
regions, identical. The nucleotide modifications present in a variant nucleic acid may be
silent, which means that they do not alter the amino acid sequences encoded by said
30 variant nucleic acid.

However, the changes in nucleotides in a variant nucleic acid may also result in substitutions, additions or deletions in the polypeptide encoded by the variant nucleic acid in relation to the polypeptides encoded by the reference nucleic acid. In addition, nucleotide modifications in the coding regions may produce conservative or non-conservative substitutions in the amino acid sequence of the polypeptide.

Preferably, the variant nucleic acids according to the invention encode polypeptides that substantially conserve the same function or biological activity as the polypeptide of the reference nucleic acid or alternatively the capacity to be recognized by antibodies directed against the polypeptides encoded by the initial reference nucleic acid.

Some variant nucleic acids will thus encode mutated forms of the polypeptides whose systematic study will make it possible to deduce structure-activity relationships of the proteins in question. Knowledge of these variants in relation to the disease studied is essential since it makes it possible to understand the molecular cause of the pathology.

AFragment@ will be understood to mean a nucleotide sequence of reduced length relative to the reference nucleic acid and comprising, over the common portion, a nucleotide sequence identical to the reference nucleic acid. Such a nucleic acid Afragment@ according to the invention may be, where appropriate, included in a larger polynucleotide of which it is a constituent. Such fragments comprise, or alternatively consist of, oligonucleotides ranging in length from 8, 10, 12, 15, 18, 20 to 25, 30, 40, 50, 70, 80, 100, 200, 500, 1000 or 1500 consecutive nucleotides of a nucleic acid according to the invention.

25

A "nucleic acid molecule" refers to the phosphate ester polymeric form of ribonucleosides (adenosine, guanosine, uridine or cytidine; "RNA molecules") or deoxyribonucleosides (deoxyadenosine, deoxyguanosine, deoxythymidine, or deoxycytidine; "DNA molecules"), or any phosphoester analogs thereof, such as phosphorothioates and thioesters, in either single stranded form, or a double-stranded helix. Double stranded DNA-DNA, DNA-RNA and RNA-RNA helices are possible. The term nucleic acid molecule, and in particular DNA or RNA molecule, refers only to

the primary and secondary structure of the molecule, and does not limit it to any particular tertiary forms. Thus, this term includes double-stranded DNA found, *inter alia*, in linear or circular DNA molecules (*e.g.*, restriction fragments), plasmids, and chromosomes. In discussing the structure of particular double-stranded DNA molecules, sequences may be described herein according to the normal convention of giving only the sequence in the 5' to 3' direction along the non-transcribed strand of DNA (*i.e.*, the strand having a sequence homologous to the mRNA). A "recombinant DNA molecule" is a DNA molecule that has undergone a molecular biological manipulation.

A nucleic acid molecule is "hybridizable" to another nucleic acid molecule, such as a cDNA, genomic DNA, or RNA, when a single stranded form of the nucleic acid molecule can anneal to the other nucleic acid molecule under the appropriate conditions of temperature and solution ionic strength (*see* Sambrook et al., *Molecular cloning: a laboratory manual*. 2ed. Cold Spring Harbor Laboratory, Cold spring Harbor, New York. (1989)).

The conditions of temperature and ionic strength determine the "stringency" of the hybridization. For preliminary screening for homologous nucleic acids, low stringency hybridization conditions, corresponding to a T_m of 55°, can be used, *e.g.*, 5x SSC, 0.1% SDS, 0.25% milk, and no formamide; or 30% formamide, 5x SSC, 0.5% SDS). Moderate stringency hybridization conditions correspond to a higher T_m , *e.g.*, 40% formamide, with 5x or 6x SCC. High stringency hybridization conditions correspond to the highest T_m , *e.g.*, 50% formamide, 5x or 6x SCC. Hybridization requires that the two nucleic acids contain complementary sequences, although depending on the stringency of the hybridization, mismatches between bases are possible. The appropriate stringency for hybridizing nucleic acids depends on the length of the nucleic acids and the degree of complementation, variables well known in the art. The greater the degree of similarity or homology between two nucleotide sequences, the greater the value of T_m for hybrids of nucleic acids having those sequences. The relative stability (corresponding to higher T_m) of nucleic acid hybridizations decreases in the following order: RNA:RNA, DNA:RNA, DNA:DNA. For hybrids of greater than 100 nucleotides in length, equations for calculating T_m have been derived (*see* Sambrook et al., *supra*, 9.50-0.51). For hybridization with shorter nucleic acids, *i.e.*, oligonucleotides, the position of mismatches becomes more important, and the length of

the oligonucleotide determines its specificity (*see* Sambrook et al., *supra*, 11.7-11.8). Preferably a minimum length for a hybridizable nucleic acid is at least about 10 nucleotides; preferably at least about 15 nucleotides; and more preferably the length is at least about 20 nucleotides.

5

In a specific embodiment, the term "standard hybridization conditions" refers to a T_m of 55°C, and utilizes conditions as set forth above. In a preferred embodiment, the T_m is 60°C; in a more preferred embodiment, the T_m is 65°C.

10

A High stringency hybridization conditions@ for the purposes of the present invention will be understood to mean the following conditions:

1- Membrane competition and PREHYBRIDIZATION:

- Mix: 40 l salmon sperm DNA (10 mg/ml)

+ 40 l human placental DNA (10 mg/ml)

15

- Denature for 5 minutes at 96°C, then immerse the mixture in ice.

- Remove the 2X SSC and pour 4 ml of formamide mix in the hybridization tube containing the membranes.

- Add the mixture of the two denatured DNAs.

- Incubation at 42°C for 5 to 6 hours, with rotation.

20

2- Labeled probe competition:

- Add to the labeled and purified probe 10 to 50 l Cot I DNA, depending on the quantity of repeats.

- Denature for 7 to 10 minutes at 95°C.

25

- Incubate at 65°C for 2 to 5 hours.

3- HYBRIDIZATION:

- Remove the prehybridization mix.

- Mix 40 l salmon sperm DNA + 40 l human placental DNA; denature for 5 min at 96°C, then immerse in ice.

30

- Add to the hybridization tube 4 ml of formamide mix, the mixture of the two DNAs and the denatured labeled probe/Cot I DNA .

- Incubate 15 to 20 hours at 42°C, with rotation.

4- Washes and Exposure:

- One wash at room temperature in 2X SSC, to rinse.
- 5 - Twice 5 minutes at room temperature 2X SSC and 0.1% SDS at 65°C.
- Twice 15 minutes at 65°C 1X SSC and 0.1% SDS at 65°C.
- Envelope the membranes in clear plastic wrap and expose.

The hybridization conditions described above are adapted to hybridization, under
 10 high stringency conditions, of a molecule of nucleic acid of varying length from
 20 nucleotides to several hundreds of nucleotides. It goes without saying that the
 hybridization conditions described above may be adjusted as a function of the length of
 the nucleic acid whose hybridization is sought or of the type of labeling chosen,
 according to techniques known to one skilled in the art. Suitable hybridization
 15 conditions may, for example, be adjusted according to the teaching contained in the
 manual by Hames and Higgins (1985), *supra* or in the manual by F. Ausubel et al.,
 Current Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, N.Y.
 (1989).

20 Nuclear Factor B

Nuclear factor B (NFB) comprises a family of eukaryotic transcription factors.
 The NFB transcription factors were first identified as factors which bound the enhancer
 elements in the light chain gene in murine B lymphocytes. Subsequent studies
 demonstrated NFB is found in almost all cells and regulates genes involved in tissue
 25 inflammation, cellular proliferation and cellular differentiation.

NFB comprises at least five subunits: p50, p52, p65 (RelA), c-Rel, and RelB
 which can form homo- and heterodimers in various combinations. Active forms of NFB
 are usually heterodimers comprised of p65 (RelA) and p50.

30

In most cells, NFB exists as an inactive heterodimer which is sequestered in the
 cytosol due to association with an inhibitory protein I-B. In response to stimuli such as

inflammatory stimuli, the I-B protein is phosphorylated and degraded, resulting in disassociation of the I-B-NFB complex and translocation of NFB into the nucleus. Once in the nucleus, NFB recognizes specific enhancer sites containing the NFB DNA binding motif and interacts with basal transcription factors to initiate RNA polymerase II mediated transcription in conjunction with the TATA box binding protein. As mentioned above, this results in the transcription of a wide variety of genes, particularly genes involved in immune and inflammatory responses.

As mentioned above, the present invention is based in part on the discovery of nuclear factor B-inducing factor (NFIF) proteins. These proteins are involved in induction of NFB. Discovery of these NFIF proteins provides different approaches to treating conditions that involve NFB-regulated inflammatory responses. NFB-regulated inflammatory responses are associated with a variety of disease states which include, but are not limited to, rheumatoid arthritis, atherosclerosis, autoimmune diseases, viral diseases, NSAID-induced gastropathy, neurodegenerative diseases, scrapie, sepsis, apoptosis, Crohn=s disease, renal disease, restenosis, brain injury/inflammation, Alzheimer=s disease, asthma, and improperly regulated expression of pleiotropic cytokines. In situations where it is desirable to increase induction of NFB for the purpose of resulting in an increased immune response, the NFIF proteins of the present invention can be introduced to or expressed in a patient to induce NFB. If, on the other hand, it is desirable to inhibit induction of NFB in order to inhibit or prevent an immune response, the provision according to the present invention of the genetic sequences encoding the NFIF proteins and of how to prepare these proteins can be used to identify compounds and compositions which inhibit or prevent expression of the NFIF proteins or which interact with NFIF proteins to inhibit or prevent their activity.

Nuclear Factor B Inducing Factor (NFIF) Proteins

The polypeptides and proteins of the present invention include recombinant polypeptides, natural polypeptides, or synthetic polypeptides, and can be of human, rabbit, or other animal origin.

The polypeptides can be isolated from natural sources, such as placental extracts, human plasma, or conditioned media from cultured cells using purification procedures known to one of skill in the art.

Alternatively, the polypeptides of the present invention can be prepared utilizing recombinant DNA technology which comprises combining a nucleic acid encoding the polypeptide thereof in a suitable vector, inserting the resulting vector into a suitable host cell, recovering the polypeptide produced by the resulting host cell, and purifying the polypeptide recovered.

The polypeptides are characterized by a reproducible single molecular weight and/or multiple set of molecular weights, chromatographic properties and elution profiles, amino acid composition and sequence, and biological activity.

Isolation of Nucleotide Sequences Encoding NFIF Polypeptides

The NFIF-14b and NFIF-7a embodiments of the present invention can be prepared by a variety of suitable methods known to those of skill in the art. Teachings on general molecular biology, microbiology, and recombinant DNA techniques within the skill of the art are explained fully in the literature. See, *e.g.*, Sambrook, Fritsch & Maniatis, *Molecular Cloning: A Laboratory Manual*, Second Edition (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (herein "Sambrook et al., 1989"); *DNA Cloning: A Practical Approach*, Volumes I and II (D.N. Glover ed. 1985); F.M. Ausubel et al. (eds.), *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc. (1994).

The cDNA sequence for NFIF-14b and NFIF-7a are presented in Figure 3. NFIF-7a is a splice variant of NFIF-14b. The term "splice variant" refers to a polypeptide encoded by an mRNA produced by alternative processing of the full length mRNA encoded by a gene or genes resulting in an mRNA that contains one or more deletions relative to the full length mRNA for the genes. As shown in Figure 4, relative to NFIF-14b, NFIF-7a has an internal deletion of base pairs 473 to 739. Given the information in the description herein on the DNA sequence of NFIF-14b and NFIF-7a

and the known methods in the art for obtaining cDNA, nucleotide sequences encoding NFIF-14b and NFIF-7a can be cloned readily and inserted into an appropriate vector for expression of these proteins *in vitro* or *in vivo*. For a description of methods relating to cloning cDNA and expression vectors, see Sambrook et al., 1989, *supra*. A "cloning vector" is a replicon, for example, a plasmid, phage or cosmid, to which another DNA segment may be attached so as to bring about the replication of the attached segment. A "replicon" is any genetic element (e.g., plasmid, chromosome, virus) that functions as an autonomous unit of DNA replication *in vivo*, *i.e.*, capable of replication under its own control. A cloning vector may be capable of replication in one cell type and expression in another (Ashuttle vector®). In preferred embodiments of the present invention, the cloning vector is capable of expression in a host cell and the "expression vector" is able to express NFIF at sufficient levels to effect an NFB regulated pathway in the cell.

A gene encoding NFIF-14b and NFIF-7a, whether genomic DNA or cDNA, can be isolated from a human genomic library or cDNA library. A DNA "coding sequence" is a double-stranded DNA sequence which is transcribed and translated into a polypeptide in a cell *in vitro* or *in vivo* when placed under the control of appropriate regulatory sequences. The DNA coding sequences and the appropriate regulatory sequences are preferably provided in an expression vector. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxyl) terminus. A coding sequence can include, but is not limited to, prokaryotic sequences, cDNA from eukaryotic mRNA, genomic DNA sequences from eukaryotic (e.g., mammalian) DNA, and even synthetic DNA sequences. If the coding sequence is intended for expression in a eukaryotic cell, a polyadenylation signal and transcription termination sequence will usually be located 3' to the coding sequence. Methods for obtaining a gene given the DNA sequence information presented herein are well known in the art. The DNA may be obtained by standard procedures known in the art from cloned DNA (e.g., a DNA "library"). It is obtained preferably from a cDNA library prepared from tissues with high level expression of the protein. The DNA may also be obtained by the cloning of genomic DNA, or fragments thereof, purified from the desired cell (See, for example, Sambrook

et al., 1989, *supra*; Glover, D.M. (ed.), 1985, DNA Cloning: A Practical Approach, MRL Press, Ltd., Oxford, U.K. Vol. I, II) or by chemical synthesis. Clones derived from genomic DNA may contain regulatory and intron DNA regions in addition to coding regions.

5

Methods for obtaining cDNA are well known in the art. Briefly, these methods include isolating a mixture of messenger RNA (mRNAs) from eukaryotic cells and employing a series of enzymatic reactions to synthesize double-stranded DNA copies (cDNAs) complementary to the isolated mRNAs. "Polymerase chain reaction" (PCR) refers to *in vitro* methods for amplifying specific DNA sequences using techniques well known in the art.

10

Regardless of the method used to obtain the desired cDNA, the double-stranded cDNA mixture is inserted into cloning vehicles by any one of many known techniques, depending at least in part on the particular vehicle used. Various insertion methods are discussed in Sambrook et al., 1989, *supra* and are well known in the art. A "cassette" refers to a segment of DNA that can be inserted into a vector at one or more specific restriction sites. The segment of DNA encodes a polypeptide of interest and the cassette and restriction sites are designed to ensure insertion of the cassette in the proper reading frame for transcription and translation.

15

20

Once the DNA segments are inserted into a cloning vehicle, the cloning vehicle is used to transform a suitable host. A cell has been "transfected" by exogenous or heterologous DNA when such DNA has been introduced inside the cell. A cell has been "transformed" by exogenous or heterologous DNA when the transfected DNA effects a phenotypic change. The transforming DNA can be integrated (covalently linked) into chromosomal DNA making up the genome of the cell. These cloning vehicles usually impart an antibiotic resistance trait on the host. Such hosts are generally prokaryotic cells and only a few of the host cells contain the desired cDNA. The transfected host cells constitute a gene library, providing a representative sample of the mRNAs present in the cell from which the mRNAs were isolated.

25

30

Given the sequence information on NFIF-14b and NFIF-7a provided herein, an appropriate oligonucleotide may be prepared, preferably synthesized as discussed above, and used to identify clones containing NFIF sequences. The oligonucleotide preferably includes at least about 18 nucleotides and is hybridizable to a genomic DNA molecule, a
5 cDNA molecule, or an mRNA molecule encoding NFIF. Oligonucleotides can be labeled, *e.g.*, with ^{32}P -nucleotides or nucleotides to which a label, such as biotin, has been covalently conjugated.

In one embodiment, a labeled oligonucleotide can be used as a probe to detect the
10 presence of a nucleic acid encoding NFIF. In another embodiment, oligonucleotides (one or both of which may be labeled) can be used as PCR primers, either for cloning full length or a fragment of NFIF, or to detect the presence of nucleic acids encoding NFIF. In a further embodiment, an oligonucleotide can form a triple helix with an NFIF DNA molecule.

15

Generally, oligonucleotides are prepared synthetically, preferably on a nucleic acid synthesizer. Accordingly, oligonucleotides can be prepared with non-naturally occurring phosphoester analog bonds, such as thioester bonds, etc. To identify clones containing the NFIF sequences, individual transformed or transfected cells are grown as
20 colonies on a nitrocellulose filter paper. The colonies are lysed and the DNA is bound tightly to the filter paper by heating. The filter paper is then incubated with a labeled oligonucleotide probe which is complementary to NFIF. DNA fragments with substantial homology to NFIF will hybridize to the probe.

25

As discussed above, the conditions of temperature and ionic strength determine the "stringency" of the hybridization. Hybridization requires that the two nucleic acids contain complementary sequences, although depending on the stringency of the hybridization, mismatches between bases are possible. The appropriate stringency for hybridizing nucleic acids depends on the length of the nucleic acids and the degree of
30 complementation, variables well known in the art.

The probe hybridizes with the cDNA for which it is complementary. It can be identified by autoradiography or by chemical reactions that identify the presence of the probe. The corresponding clones are characterized in order to identify one or a combination of clones which contain all of the structural information for the desired protein. The nucleic acid sequence coding for the protein of interest is isolated and reinserted into an expression vector. The expression vector brings the cloned gene under the regulatory control of specific prokaryotic or eukaryotic control elements which allow the efficient expression (transcription and translation) of the ds-cDNA.

Transcriptional and translational control sequences are DNA regulatory sequences, such as, for example, promoters, enhancers, and terminators that provide for the expression of a coding sequence in a host cell. In eukaryotic cells, polyadenylation signals are control sequences. A coding sequence is "under the control" of transcriptional and translational control sequences in a cell when RNA polymerase transcribes the coding sequence into mRNA, which is then spliced (if the coding sequence contains introns) and translated into the protein encoded by the coding sequence.

Further selection can be carried out on the basis of the properties of the gene. For example, the presence of the desired gene in a clone may be detected by assays based on the physical, chemical, or immunological properties of its expressed protein product. For example, cDNA clones, or DNA clones can be selected which produce a protein that has similar or identical properties to those of NFIF with regard to electrophoretic migration, isoelectric focusing, non-equilibrium pH gel electrophoresis, proteolytic digestion, or antigenicity.

Preparation of Variants of NFIF Polypeptides

The present invention includes within its scope allelic variants, substitution, addition and deletion mutant variants, analogs, and derivatives of NFIF and homologs from other species that have the same or homologous functional activity as NFIF. In preferred embodiments, genes having deletions or substitutions that increase the ability to induce NFB are utilized in the practice of the invention. Preparation or isolation of NFIF variants are within the scope of the present invention. Accordingly, the scope of

the present invention includes NFIF variants which are functionally active, *i.e.*, capable of exhibiting one or more functional activities associated with NFIF.

NFIF variants can be made by altering encoding nucleic acid sequences by
5 substitutions, additions or deletions that provide for functionally equivalent molecules. Preferably, NFIF embodiments are made that have enhanced or increased functional activity relative to NFIF-14b or NFIF-7a.

Due to the degeneracy of nucleotide coding sequences, other DNA sequences
10 which encode substantially the same amino acid sequence as NFIF, including an amino acid sequence that contains a single amino acid variant, can be used in the practice of the present invention. These include, but are not limited to, allelic genes, homologous genes from other species, and nucleotide sequences comprising all or portions of NFIF which are altered by the substitution of different codons that encode the same amino acid
15 residue within the sequence, thus producing a silent change. Likewise, the NFIF variants of the invention include, but are not limited to, those containing, as a primary amino acid sequence, all or part of the amino acid sequence of a NFIF protein including altered sequences in which functionally equivalent amino acid residues are substituted for residues within the sequence resulting in a conservative amino acid substitution. For
20 example, one or more amino acid residues within the sequence can be substituted by another amino acid of a similar polarity, which acts as a functional equivalent, resulting in a silent alteration. Substitutes for an amino acid within the sequence can be selected from other members of the class to which the amino acid belongs. For example, the nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline,
25 phenylalanine, tryptophan and methionine. Amino acids containing aromatic ring structures are phenylalanine, tryptophan, and tyrosine. The polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine. The positively charged (basic) amino acids include arginine, lysine and histidine. The negatively charged (acidic) amino acids include aspartic acid and glutamic acid. Such
30 alterations will not be expected to affect apparent molecular weight as determined by polyacrylamide gel electrophoresis, or isoelectric point.

Particularly preferred substitutions are:

- Lys for Arg and vice versa such that a positive charge may be maintained;
- Glu for Asp and vice versa such that a negative charge may be maintained;
- Ser for Thr such that a free -OH can be maintained; and
- 5 - Gln for Asn such that a free CONH₂ can be maintained.

Amino acid substitutions may also be introduced to substitute an amino acid with a particularly preferable property. For example, a Cys may be introduced as a potential site for disulfide bridges with another Cys. A His may be introduced as a particularly
10 "catalytic" site (*i.e.*, His can act as an acid or base and is the most common amino acid in biochemical catalysis). Pro may be introduced because of its particularly planar structure, which induces -turns in the protein's structure.

The genes encoding NFIF variants of the invention can be produced by various
15 methods known in the art. The manipulations which result in their production can occur at the gene or protein level. For example, the cloned NFIF gene sequence can be modified by any of numerous strategies known in the art (Sambrook et al., 1989, *supra*).

The sequence can be cleaved at appropriate sites with restriction endonuclease(s), followed by further enzymatic modification if desired, isolated, and ligated *in vitro*. In
20 the production of the gene encoding a NFIF embodiment, care should be taken to ensure that the modified gene remains within the same translational reading frame as the NFIF gene, uninterrupted by translational stop signals, in the gene region where the desired activity is encoded.

25 Additionally, the NFIF-encoding nucleic acid sequence can be mutated *in vitro* or *in vivo* to create and/or destroy translation, initiation, and/or termination sequences, or to create variations in coding regions and/or form new restriction endonuclease sites or destroy preexisting ones, to facilitate further *in vitro* modification. Preferably, such mutations enhance the functional activity of the mutated NFIF gene product. Any
30 technique for mutagenesis known in the art can be used, for example, *in vitro* site-directed mutagenesis (Hutchinson, C., et al., 1978, *J. Biol. Chem.* 253:6551; Zoller and Smith, 1984, *DNA* 3:479-488; Oliphant et al., 1986, *Gene* 44:177; Hutchinson et al.,

1986, *Proc. Natl. Acad. Sci. U.S.A.* 83:710), and use of ATAB@ linkers (Pharmacia), etc. PCR techniques are preferred for site directed mutagenesis (see Higuchi, 1989, "Using PCR to Engineer DNA", in PCR Technology: Principles and Applications for DNA Amplification, H. Erlich, ed., Stockton Press, Chapter 6, pp. 61-70).

5

Polypeptides Based on NFIF

A therapeutically useful variant of NFIF may be identified by a variety of methods, including *in vitro* assays that can be used to identify the level of expression of an NFB regulated gene or a gene whose expression is regulated by NFB regulatory elements in the presence of the NFIF variant. One method for identifying polypeptides which are able to enhance (or inhibit) NFB induction is to use a reporter gene system. These systems utilize reporter gene expression vectors which include a cloning site into which a given promoter may be cloned upstream of a Areporter gene@ which can be easily detected and quantified. One of skill in the art could readily identify and subclone the promoter for NFB as well as other control sequences into a commercially available reporter gene expression vector. The expression vector is transferred into host cells and the cells are exposed to an NFIF variant (or a putative inhibitor or enhancer molecule) to determine the effect on expression of the reporter gene product. In particular, the cells are assayed for the presence of the reporter gene product directly by measuring the amount of reporter mRNA, the reporter protein itself or the enzymatic activity of the reporter protein. Ideally, the reporter gene is not endogenously expressed in the cell type of interest and lends itself to sensitive, quantitative and rapid assays. A variety of reporter assay constructs are commercially available and several reporter genes and assays have been developed and can be readily prepared by those of skill in the art. The most popular systems for monitoring genetic activity in eukaryotic cells include chloramphenicol acetyltransferase (CAT), -galactosidase, firefly luciferase, growth hormone (GH), -glucuronidase (GUS), alkaline phosphatase (AP), green fluorescent protein (GFP) and *Renilla* luciferase. Reporter assay constructs can be purchased from a variety of sources including Promega and Invitrogen.

30

As mentioned above, reporter gene activity can be detected by assaying for the reporter mRNA or the reporter protein. The reporter mRNA can be detected by northern blot analysis, ribonuclease protection assays or RT-PCR. While these assays are more direct than measuring protein expression, many assays have been developed to measure the presence of the reporter protein rather than the mRNA present in a cell. Reporter proteins can be assayed by spectrophotometry or by detecting enzymatic activity. Reporter protein levels may also be measured with antibody-based assays. In general, the enzymatic assays are very sensitive and are a preferred method of monitoring reporter gene expression. A preferred commercially available NFB reporter gene construct is the pNFB-Luc (luciferase) reporter gene vector available from Stratagene. An example of how to utilize this reporter system to quantify NFIF protein activity is provided in Example 4.

Experiments of the type discussed hereinabove can be utilized to determine how well a given disease state can be treated using the compositions of the present invention.

The discussion which follows relates to the manipulation and expression of DNA encoding the polypeptides of the present invention.

Expression Vectors Encoding NFIF Polypeptides

The identified and isolated DNA sequence can be inserted into an appropriate cloning/expression vector (hereinafter "vector") to facilitate modifications to the sequence or expression of the protein. These vectors typically include multiple cloning sites, promoters, sequences which facilitate replication in a host cell and selection markers.

Any suitable vector can be used. There are many known in the art. Examples of vectors that can be used include, for example, plasmids or modified viruses. The vector is typically compatible with a given host cell into which the vector is introduced to facilitate replication of the vector and expression of the encoded proteins. The insertion of a DNA sequence into a given vector can, for example, be accomplished by ligating the DNA fragment into a cloning vector which has complementary cohesive termini.

However, if the complementary restriction sites used to cut the DNA are not present in the cloning vector, the ends of the DNA molecules may be enzymatically modified. Alternatively, any site desired may be produced by ligating nucleotide sequences (linkers) onto the DNA termini; the ligated linkers may comprise specific chemically synthesized oligonucleotides encoding restriction endonuclease recognition sequences. Useful vectors may consist of segments of chromosomal, non-chromosomal and synthetic DNA sequences. Examples of specific vectors useful in the practice of the present invention are *E. coli* bacteriophages, for example, lambda derivatives, or plasmids, for example, pBR322 derivatives or pUC plasmid derivatives, *e.g.*, pmal-c, pFLAG, derivatives of SV40 and known bacterial plasmids, *e.g.*, *E. coli* plasmids col El, pCR1, pMal-C2, pET, pGEX (Smith *et al.*, 1988, *Gene* 67:31-40), pMB9 and their derivatives, plasmids such as RP4; phage DNAs, *e.g.*, the numerous derivatives of phage 1, *e.g.*, NM989, and other phage DNA, *e.g.*, M13 and filamentous single stranded phage DNA; yeast vectors such as the 2 m plasmid or derivatives thereof; vectors useful in eukaryotic cells, for example, vectors useful in insect cells, such as baculovirus vectors, vectors useful in mammalian cells; vectors derived from combinations of plasmids and phage DNAs, plasmids that have been modified to employ phage DNA or other expression control sequences; and the like.

Examples of yeast vectors that can be used according to the invention are the non-fusion pYES2 vector (Invitrogen) or the fusion pYESHisA, B, C (Invitrogen).

Baculovirus vectors that can be used in the practice of the invention include a variety of vectors, including both non-fusion transfer vectors, for example, pVL941 (Summers), pVL1393 (Invitrogen), pVL1392 (Summers and Invitrogen), and pBlueBacIII (Invitrogen), and fusion transfer vectors, for example, pAc700 (Summers), pAc701 and pAc702, pAc360 (Invitrogen), and pBlueBacHisA, B, C (Invitrogen) can be used.

Mammalian vectors contemplated for use in the invention include, for example, vectors with inducible promoters, for example, the dihydrofolate reductase (DHFR) promoter, *e.g.*, any expression vector with a *DHFR* expression vector, or a

DHFR/methotrexate co-amplification vector, for example, pED (*see* Kaufman, *Current Protocols in Molecular Biology*, 16.12 (1991)). Alternatively, a glutamine synthetase/methionine sulfoximine co-amplification vector, for example, pEE14 (Celltech). In another embodiment, a vector that directs episomal expression under control of Epstein Barr Virus (EBV) can be used, for example, pREP4 (Invitrogen), pCEP4 (Invitrogen), pMEP4 (Invitrogen), pREP8 (Invitrogen), pREP9 (Invitrogen), and pEBVHis (Invitrogen). Selectable mammalian expression vectors for use in the invention include pRc/CMV (Invitrogen), pRc/RSV (Invitrogen), pcDNA3 (Invitrogen) and others. Vaccinia virus mammalian expression vectors (*see*, Kaufman, 1991, *supra*) for use according to the invention include but are not limited to pSC11, pMJ601, and pTKgptF1S.

A variety of methods may be used to confirm that the desired DNA sequence encoding NFIF-14b, NFIF-7a, or another NFIF variant, has been cloned into a vector. In general, one or more of the following approaches is used: (a) PCR amplification of the desired plasmid DNA or specific mRNA, (b) nucleic acid hybridization, (c) presence or absence of selection marker gene functions, (d) analyses with appropriate restriction endonucleases, and (e) expression of inserted sequences. In the first approach, the nucleic acids can be amplified by PCR to provide for detection of the amplified product. In the second approach, the presence of a foreign gene inserted in an expression vector can be detected by nucleic acid hybridization using probes comprising sequences that are homologous to an inserted marker gene. In the third approach, the recombinant vector/host system can be identified and selected based upon the presence or absence of certain "selection marker" gene functions (*e.g.*, -galactosidase activity, thymidine kinase activity, resistance to antibiotics, transformation phenotype, occlusion body formation in baculovirus, etc.) caused by the insertion of foreign genes in the vector. In another example, if the nucleic acid encoding NFIF is inserted within the "selection marker" gene sequence of the vector, recombinants containing the NFIF insert can be identified by the absence of the selection marker gene function. In the fourth approach, recombinant expression vectors are identified by digestion with appropriate restriction enzymes. In the fifth approach, recombinant expression vectors can be identified by assaying for the activity, biochemical, or immunological characteristics of the gene

product expressed by the recombinant, provided that the expressed protein assumes a functionally active conformation.

Promoters

5 The nucleotide sequence coding for NFIF-14b or NFIF-7a or a NFIF variant thereof can be inserted into an expression vector which contains the necessary elements for the transcription and translation of the inserted protein-coding sequence. Such elements are termed herein a "promoter."

10 A promoter is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. For purposes of defining the present invention, the promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements necessary to initiate transcription at
15 levels detectable above background. Within the promoter sequence will be found a transcription initiation site (conveniently defined for example, by mapping with nuclease S1), as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase.

20 The nucleic acid encoding the polypeptides of the invention is operationally associated with a promoter in an expression vector of the invention. Both cDNA and genomic sequences can be cloned and expressed under control of such regulatory sequences. An expression vector also preferably includes a replication origin. The necessary transcriptional and translational signals can be provided on a recombinant
25 expression vector or they may be supplied by the native gene encoding NFIF and/or its flanking regions. Any of the methods previously described for the insertion of DNA fragments into a cloning vector may be used to construct expression vectors containing a gene consisting of appropriate transcriptional/translational control signals and the protein coding sequences. These methods may include *in vitro* recombinant DNA and synthetic
30 techniques and *in vivo* recombination (genetic recombination).

Expression may be controlled by any promoter/enhancer element known in the art, but these regulatory elements must be functional in the host selected for expression.

Examples of promoters which may be used to control NFIF gene expression include the SV40 early promoter region (Benoist and Chambon, 1981, *Nature* 290:304-310), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto, et al., 1980, *Cell* 22:787-797), the herpes thymidine kinase promoter (Wagner et al., 1981, *Proc. Natl. Acad. Sci. U.S.A.* 78:1441-1445), the regulatory sequences of the metallothionein gene (Brinster et al., 1982, *Nature* 296:39-42); prokaryotic expression vectors for example, the -lactamase promoter (Villa-Kamaroff, et al., 1978, *Proc. Natl. Acad. Sci. U.S.A.*, 75:3727-3731), or the *tac* promoter (DeBoer, et al., 1983, *Proc. Natl. Acad. Sci. U.S.A.*, 80:21-25); promoter elements from yeast or other fungi, for example, the Gal 4 promoter, the ADC (alcohol dehydrogenase) promoter, PGK (phosphoglycerol kinase) promoter, alkaline phosphatase promoter; and the animal transcriptional control regions, which exhibit tissue specificity and have been utilized in transgenic animals: elastase I gene control region which is active in pancreatic acinar cells (Swift et al., 1984, *Cell* 38:639-646; Ornitz et al., 1986, Cold Spring Harbor *Symp. Quant. Biol.*, 50:399-409; MacDonald, 1987, *Hepatology* 7:425-515); insulin gene control region which is active in pancreatic beta cells (Hanahan, 1985, *Nature* 315:115-122), immunoglobulin gene control region which is active in lymphoid cells (Grosschedl et al., 1984, *Cell* 38:647-658; Adames et al., 1985, *Nature* 318:533-538; Alexander et al., 1987, *Mol. Cell. Biol.*, 7:1436-1444), mouse mammary tumor virus control region which is active in testicular, breast, lymphoid and mast cells (Leder et al., 1986, *Cell* 45:485-495), albumin gene control region which is active in liver (Pinkert et al., 1987, *Genes and Devel.* 1:268-276), alpha-fetoprotein gene control region which is active in liver (Krumlauf et al., 1985, *Mol. Cell. Biol.*, 5:1639-1648; Hammer et al., 1987, *Science* 235:53-58), alpha 1-antitrypsin gene control region which is active in the liver (Kelsey et al., 1987, *Genes and Devel.*, 1:161-171), beta-globin gene control region which is active in myeloid cells (Mogam et al., 1985, *Nature* 315:338-340; Kollias et al., 1986, *Cell* 46:89-94), myelin basic protein gene control region which is active in oligodendrocyte cells in the brain (Readhead et al., 1987, *Cell* 48:703-712), myosin light chain-2 gene control region which is active in skeletal muscle (Sani, 1985,

Nature 314:283-286), and gonadotropic releasing hormone gene control region which is active in the hypothalamus (Mason et al., 1986, *Science* 234:1372-1378).

A preferred promoter used in the expression vector constructs discussed
5 hereinbelow is the cytomegalovirus (CMV) promoter which is capable of providing high level expression in a variety of mammalian cell lines.

Introduction of Vectors into Host Cells

Vectors can be introduced into host cells by any suitable method, including, *e.g.*,
10 transfection, electroporation, microinjection, transduction, cell fusion, DEAE dextran, calcium phosphate precipitation, lipofection (lysosome fusion), use of a gene gun, or a DNA vector transporter (see, *e.g.*, Wu et al., 1992, *J. Biol. Chem.* 267:963-967; Wu and Wu, 1988, *J. Biol. Chem.* 263:14621-14624; Hartmut et al., Canadian Patent Application No. 2,012,311, filed March 15, 1990) so that many copies of the gene
15 sequence are generated. In a preferred method, cells are transfected *in vitro* utilizing Lipfectamine7 available from Gibco-BRL. Preferably, the cloned gene is contained on a shuttle vector plasmid, which provides for expansion in a cloning cell, *e.g.*, *E. coli*, and facilitates purification for subsequent insertion into an appropriate expression cell line. For example, a shuttle vector, which is a vector that can replicate in more than one type
20 of organism, can be prepared for replication in both *E. coli* and *Saccharomyces cerevisiae* by linking sequences from an *E. coli* plasmid with sequences from the yeast 2 m plasmid.

Host Cell Systems

25 Host cell systems include mammalian host cell systems, insect host cell systems and microorganisms such as yeast or bacteria. Depending on the host cell system utilized, any one of a number of suitable transcription and translation elements may be used.

30 In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences or modifies and processes the gene product in the specific fashion desired. Different host cells have characteristic and specific mechanisms for the

translational and post-translational processing and modification of proteins. Appropriate cell lines or host systems can be chosen to ensure the desired modification and processing of the foreign protein expressed. Expression in yeast can produce a biologically active product. Expression in eukaryotic cells can increase the likelihood of "native" folding. Moreover, expression in mammalian cells can provide a tool for reconstituting, or constituting, NFIF-inhibiting activity. Furthermore, different vector/host expression systems may affect processing reactions, such as proteolytic cleavages, to a different extent. Expression vectors of the invention can be used, as pointed out above, both to transfect cells for screening or biological testing of modulators of NFIF activity.

Examples of acceptable mammalian host cells are HEK 293 cells and COS-7 cells.

A recombinant NFIF-14b, NFIF-7a or NFIF variant of the invention may be expressed chromosomally, after integration of the coding sequence by recombination. In this regard, any of a number of amplification systems may be used to achieve high levels of stable gene expression (*See* Sambrook et al., 1989, *supra*).

The cell into which the recombinant vector comprising the nucleic acid encoding NFIF is introduced is cultured in an appropriate cell culture medium under conditions that provide for expression of an NFIF polypeptide by the cell.

Once a suitable host system and growth conditions are established, recombinant expression vectors can be propagated and prepared in quantity. Soluble forms of the protein can be obtained by collecting culture fluid, or solubilizing inclusion bodies, *e.g.*, by treatment with detergent, and if desired sonication or other mechanical processes, as described above. The solubilized or soluble protein can be isolated using various techniques, including polyacrylamide gel electrophoresis (PAGE), isoelectric focusing, 2-dimensional gel electrophoresis, chromatography (*e.g.*, ion exchange, affinity, immunoaffinity, and sizing column chromatography), centrifugation, differential

solubility, immunoprecipitation, or by any other standard technique for the purification of proteins.

As discussed above, a Avector@ is any means for the transfer of a nucleic acid
5 according to the invention into a host cell. Preferred vectors are viral vectors, for example, retroviruses, herpes viruses, adenoviruses, and adeno-associated viruses. Thus, a gene encoding a protein or polypeptide domain fragment of the present invention is introduced *in vivo*, *ex vivo*, or *in vitro* using a viral vector or through direct introduction of DNA. Expression in targeted tissues can be effected by targeting the
10 transgenic vector to specific cells, such as with a viral vector or a receptor ligand, or by using a tissue-specific promoter, or both.

The discussion which follows outlines various viral and non-viral systems that may be used to introduce DNA into a host cell *in vivo* or *in vitro*.

15

Viral Vector Systems

The NFIF polypeptides, as well as the antisense nucleic acids, ribozymes and antibodies discussed hereinbelow may be prepared *in vitro* or *ex vivo* or may be designed
20 to be expressed *in vivo* in a patient using an appropriate expression system introduced via a viral vector system.

Viral vectors commonly used for *in vivo* or *ex vivo* targeting and therapy procedures are DNA-based vectors and retroviral vectors. Methods for constructing and
25 using viral vectors are known in the art [*see, e.g.*, Miller and Rosman, *BioTechniques* 7:980-990 (1992)]. Preferably, the viral vectors are replication defective, that is, they are unable to replicate autonomously in the target cell. In general, the genome of the replication defective viral vectors which are used within the scope of the present invention lack at least one region which is necessary for the replication of the virus in
30 the infected cell. These regions can either be eliminated (in whole or in part), or be rendered non-functional by any technique known to a person skilled in the art. These techniques include the total removal, substitution (by other sequences, in particular by

the inserted nucleic acid), partial deletion or addition of one or more bases to an essential (for replication) region. Such techniques may be performed *in vitro* (on the isolated DNA) or *in situ*, using the techniques of genetic manipulation or by treatment with mutagenic agents. Preferably, the replication defective virus retains the sequences
 5 of its genome which are necessary for encapsulating the viral particles.

DNA viral vectors include an attenuated or defective DNA virus, such as but not limited to herpes simplex virus (HSV), papillomavirus, Epstein Barr virus (EBV), adenovirus, adeno-associated virus (AAV), vaccinia virus, and the like. Defective
 10 viruses, which entirely or almost entirely lack viral genes, are preferred. Defective virus is not replication competent after introduction into a cell, and thus does not lead to a productive viral infection. Use of defective viral vectors allows for administration to cells in a specific, localized area, without concern that the vector can infect other cells. Thus, a specific tissue can be specifically targeted. Examples of particular vectors
 15 include, but are not limited to, a defective herpes virus 1 (HSV1) vector (Kaplitt et al., *Molec. Cell. Neurosci.* 2:320-330 (1991)), defective herpes virus vector lacking a glycoprotein L gene (Patent Publication RD 371005 A), or other defective herpes virus vectors (International Patent Publication No. WO 94/21807, published September 29, 1994; International Patent Publication No. WO 92/05263, published April 2, 1994); an
 20 attenuated adenovirus vector, such as the vector described by Stratford-Perricaudet et al. (*J. Clin. Invest.* 90:626-630 (1992); see also La Salle et al., *Science* 259:988-990 (1993)); and a defective adeno-associated virus vector (Samulski et al., *J. Virol.* 61:3096-3101 (1987); Samulski et al., *J. Virol.* 63:3822-3828 (1989); Lebkowski et al., *Mol. Cell. Biol.* 8:3988-3996 (1988))]

25 Preferably, for *in vivo* administration, an appropriate immunosuppressive treatment is employed in conjunction with the viral vector, *e.g.*, adenovirus vector, to avoid immuno-deactivation of the viral vector and transfected cells. For example, immunosuppressive cytokines, such as interleukin-12 (IL-12), interferon- (IFN-), or
 30 anti-CD4 antibody, can be administered to block humoral or cellular immune responses to the viral vectors. In addition, it is advantageous to employ a viral vector that is engineered to express a minimal number of antigens.

Naturally, the invention contemplates delivery of a vector that will express a therapeutically effective amount of NFIF for gene therapy applications. The phrase "therapeutically effective amount" is used herein to mean an amount sufficient to cause an improvement in a clinically significant condition in the host.

Certain viral vector systems are well developed in the art and are suited to the treatment methods of the present invention.

Adenovirus Vector Systems

In a preferred embodiment, the vector is an adenovirus vector. Adenoviruses are eukaryotic DNA viruses that can be modified to efficiently deliver a nucleic acid of the invention to a variety of cell types. Various serotypes of adenovirus exist. Of these serotypes, preference is given, within the scope of the present invention, to using type 2 or type 5 human adenoviruses (Ad 2 or Ad 5) or adenoviruses of animal origin (see WO94/26914). Those adenoviruses of animal origin which can be used within the scope of the present invention include adenoviruses of canine, bovine, murine (example: Mav1, Beard et al., *Virology* 75 (1990) 81), bovine, porcine, avian, and simian (example: SAV) origin.

Preferably, the replication defective adenoviral vectors of the invention comprise the ITRs, an encapsidation sequence and the nucleic acid of interest. Still more preferably, at least the E1 region of the adenoviral vector is non-functional. The deletion in the E1 region preferably extends from nucleotides 455 to 3329 in the sequence of the Ad5 adenovirus (PvuII-BglII fragment) or 382 to 3446 (HinfII-Sau3A fragment). Other regions may also be modified, in particular the E3 region (WO95/02697), the E2 region (WO94/28938), the E4 region (WO94/28152, WO94/12649 and WO95/02697), or in any of the late genes L1-L5.

In a preferred embodiment, the adenoviral vector has a deletion in the E1 region (Ad 1.0). Examples of E1-deleted adenoviruses are disclosed in EP 185,573, the contents of which are incorporated herein by reference. In another preferred

embodiment, the adenoviral vector has a deletion in the E1 and E4 regions (Ad 3.0). Examples of E1/E4-deleted adenoviruses are disclosed in WO95/02697 and WO96/22378, the contents of which are incorporated herein by reference. In still another preferred embodiment, the adenoviral vector has a deletion in the E1 region into which the E4 region and the nucleic acid sequence are inserted (see FR94 13355, the contents of which are incorporated herein by reference).

The replication defective recombinant adenoviruses according to the invention can be prepared by any technique known to the person skilled in the art (Levrero et al., *Gene* 101 (1991) 195, EP 185 573; Graham, *EMBO J.* 3 (1984) 2917). In particular, they can be prepared by homologous recombination between an adenovirus and a plasmid which carries, inter alia, the DNA sequence of interest. The homologous recombination is effected following cotransfection of the adenovirus and plasmid into an appropriate cell line. The cell line which is employed should preferably (i) be transformable by the said elements, and (ii) contain the sequences which are able to complement the part of the genome of the replication defective adenovirus, preferably in integrated form in order to avoid the risks of recombination. Examples of cell lines which may be used are the human embryonic kidney cell line 293 (Graham et al., *J. Gen. Virol.* 36 (1977) 59) which contains the left-hand portion of the genome of an Ad5 adenovirus (12%) integrated into its genome, and cell lines which are able to complement the E1 and E4 functions, as described in applications WO94/26914 and WO95/02697. Recombinant adenoviruses are recovered and purified using standard molecular biological techniques, which are well known to one of ordinary skill in the art.

Adeno-Associated Virus Vector Systems

The adeno-associated viruses (AAV) are DNA viruses of relatively small size which can integrate, in a stable and site-specific manner, into the genome of the cells which they infect. They are able to infect a wide spectrum of cells without inducing any effects on cellular growth, morphology or differentiation, and they do not appear to be involved in human pathologies. The AAV genome has been cloned, sequenced and characterised. It encompasses approximately 4700 bases and contains an inverted

terminal repeat (ITR) region of approximately 145 bases at each end, which serves as an origin of replication for the virus. The remainder of the genome is divided into two essential regions which carry the encapsulation functions: the left-hand part of the genome, which contains the rep gene involved in viral replication and expression of the viral genes; and the right-hand part of the genome, which contains the cap gene encoding the capsid proteins of the virus.

The use of vectors derived from the AAVs for transferring genes *in vitro* and *in vivo* has been described (see WO 91/18088; WO 93/09239; US 4,797,368, US 5,139,941, EP 488 528). These publications describe various AAV-derived constructs in which the rep and/or cap genes are deleted and replaced by a gene of interest, and the use of these constructs for transferring the gene of interest *in vitro* (into cultured cells) or *in vivo*, (directly into an organism). The replication defective recombinant AAVs according to the invention can be prepared by cotransfecting a plasmid containing the nucleic acid sequence of interest flanked by two AAV inverted terminal repeat (ITR) regions, and a plasmid carrying the AAV encapsulation genes (rep and cap genes), into a cell line which is infected with a human helper virus (for example an adenovirus). The AAV recombinants which are produced are then purified by standard techniques.

The invention also relates, therefore, to an AAV-derived recombinant virus whose genome encompasses a sequence encoding a nucleic acid encoding NFIF or its variants flanked by the AAV ITRs. The invention also relates to a plasmid encompassing a sequence encoding a nucleic acid encoding NFIF or its variants flanked by two ITRs from an AAV. Such a plasmid can be used as it is for transferring the nucleic acid sequence, with the plasmid, where appropriate, being incorporated into a liposomal vector (pseudo-virus).

Retrovirus Vector Systems

In another embodiment the gene can be introduced in a retroviral vector, *e.g.*, as described in Anderson et al., U.S. Patent No. 5,399,346; Mann et al., 1983, *Cell* 33:153; Temin et al., U.S. Patent No. 4,650,764; Temin et al., U.S. Patent No. 4,980,289; Markowitz et al., 1988, *J. Virol.* 62:1120; Temin et al., U.S. Patent No.

5,124,263; EP 453242, EP178220; Bernstein et al. *Genet. Eng.* 7 (1985) 235; McCormick, *BioTechnology* 3 (1985) 689; International Patent Publication No. WO 95/07358, published March 16, 1995, by Dougherty et al.; and Kuo et al., 1993, *Blood* 82:845. The retroviruses are integrating viruses which infect dividing cells. The
5 retrovirus genome includes two LTRs, an encapsulation sequence and three coding regions (gag, pol and env). In recombinant retroviral vectors, the *gag*, *pol* and *env* genes are generally deleted, in whole or in part, and replaced with a heterologous nucleic acid sequence of interest. These vectors can be constructed from different types of retrovirus, such as, HIV, MoMuLV ("murine Moloney leukaemia virus" MSV
10 ("murine Moloney sarcoma virus"), HaSV ("Harvey sarcoma virus"); SNV ("spleen necrosis virus"); RSV ("Rous sarcoma virus") and Friend virus. Defective retroviral vectors are disclosed in WO95/02697.

In general, in order to construct recombinant retroviruses containing a nucleic
15 acid sequence, a plasmid is constructed which contains the LTRs, the encapsulation sequence and the coding sequence. This construct is used to transfect a packaging cell line, which cell line is able to supply in trans the retroviral functions which are deficient in the plasmid. In general, the packaging cell lines are thus able to express the *gag*, *pol* and *env* genes. Such packaging cell lines have been described in the prior art, in
20 particular the cell line PA317 (US 4,861,719); the PsiCRIP cell line (WO90/02806) and the GP+envAm-12 cell line (WO89/07150). In addition, the recombinant retroviral vectors can contain modifications within the LTRs for suppressing transcriptional activity as well as extensive encapsulation sequences which may include a part of the *gag* gene (Bender et al., *J. Virol.* 61 (1987) 1639). Recombinant retroviral vectors are
25 purified by standard techniques known to those having ordinary skill in the art.

Retroviral vectors can be constructed to function as infections particles or to undergo a single round of transfection. In the former case, the virus is modified to retain all of its genes except for those responsible for oncogenic transformation
30 properties, and to express the heterologous gene. Non-infectious viral vectors are prepared to destroy the viral packaging signal, but retain the structural genes required to package the co-introduced virus engineered to contain the heterologous gene and the

packaging signals. Thus, the viral particles that are produced are not capable of producing additional virus.

5 Non-Viral Systems

Certain non-viral systems have been used in the art and can facilitate introduction of DNA encoding the NFIF polypeptides, antisense nucleic acids, ribozymes and antibodies.

10 Lipofection Delivery Systems

A vector can be introduced *in vivo* by lipofection. For the past decade, there has been increasing use of liposomes for encapsulation and transfection of nucleic acids *in vitro*. Synthetic cationic lipids designed to limit the difficulties and dangers encountered with liposome mediated transfection can be used to prepare liposomes for *in vivo* transfection of a gene encoding a marker (Felgner, et. al., *Proc. Natl. Acad. Sci. U.S.A.* 84:7413-7417 (1987); see Mackey, et al., *Proc. Natl. Acad. Sci. U.S.A.* 85:8027-8031 (1988); Ulmer et al., *Science* 259:1745-1748 (1993)). The use of cationic lipids may promote encapsulation of negatively charged nucleic acids, and also promote fusion with negatively charged cell membranes (Felgner and Ringold, *Science* 337:387-388 (1989)).

20 Particularly useful lipid compounds and compositions for transfer of nucleic acids are described in International Patent Publications WO95/18863 and WO96/17823, and in U.S. Patent No. 5,459,127. The use of lipofection to introduce exogenous genes into the specific organs *in vivo* has certain practical advantages. Molecular targeting of liposomes to specific cells represents one area of benefit. It is clear that directing

25 transfection to particular cell types would be particularly advantageous in a tissue with cellular heterogeneity, for example, pancreas, liver, kidney, and the brain. Lipids may be chemically coupled to other molecules for the purpose of targeting (see Mackey, et. al., *supra*). Targeted peptides, *e.g.*, hormones or neurotransmitters, and proteins for example, antibodies, or non-peptide molecules could be coupled to liposomes

30 chemically.

Other molecules are also useful for facilitating transfection of a nucleic acid *in vivo*, for example, a cationic oligopeptide (*e.g.*, International Patent Publication WO95/21931), peptides derived from DNA binding proteins (*e.g.*, International Patent Publication WO96/25508), or a cationic polymer (*e.g.*, International Patent Publication
5 WO95/21931).

Naked DNA Delivery Systems

It is also possible to introduce the vector *in vivo* as a naked DNA plasmid (see U.S. Patents 5,693,622, 5,589,466 and 5,580,859). Naked DNA vectors for gene
10 therapy can be introduced into the desired host cells by methods known in the art, *e.g.*, transfection, electroporation, microinjection, transduction, cell fusion, DEAE dextran, calcium phosphate precipitation, use of a gene gun, or use of a DNA vector transporter (*see, e.g.*, Wu et al., *J. Biol. Chem.* 267:963-967 (1992); Wu and Wu, *J. Biol. Chem.* 263:14621-14624 (1988); Hartmut et al., Canadian Patent Application No. 2,012,311,
15 filed March 15, 1990; Williams et al., *Proc. Natl. Acad. Sci. USA* 88:2726-2730 (1991)). Receptor-mediated DNA delivery approaches can also be used (Curiel et al., *Hum. Gene Ther.* 3:147-154 (1992); Wu and Wu, *J. Biol. Chem.* 262:4429-4432 (1987)).

Uses for NFIF Polypeptides

As mentioned above, in certain situations it is desirable to induce NFB in order to initiate or increase the extent of an immune response. In order to induce NFB, the NFIF polypeptides of the present invention can be introduced into the body of a patient by a variety of methods.

25 The methods used to introduce NFIF into a patient's body include direct administration of purified NFIF polypeptides or introduction of nucleic acid encoding NFIF polypeptides in expression vectors which express the polypeptides in the patient's body.

In embodiments involving direct administration of NFIF polypeptides, the NFIF polypeptides are prepared using the host cell expression systems such as those described above. The polypeptides are purified using conventional purification methods and then combined with a suitable biologically-compatible solution as described in detail below.

5 The solution containing the polypeptide is introduced into the patient by topical, oral, parenteral, intranasal, subcutaneous or intraocular routes. Once in the body, the polypeptide is able to exert its effect, inducing NFB and thereby resulting in an increase in the activity of NFB-regulated pathways, including immune responses.

10 In embodiments involving administration of nucleic acids encoding NFIF polypeptides, viral or non-viral systems utilizing expression vectors capable of expressing NFIF polypeptides can be introduced into the patient's body. Any of the viral or non-viral transfection systems described above may be used. The viral and non-viral vector systems are also combined with an appropriate biologically-compatible
15 solution to facilitate their introduction into the body. Once a viral or non-viral vector is introduced into the body, the NFIF polypeptide encoding nucleic acid may integrate into the host's genome, providing stable, long-term expression of NFIF polypeptides which exert their effect as described above. Transient expression may be provided by systems that introduce the nucleic acid into cells, but do not integrate into the genome.

20

The NFIF-14b and NFIF-7a polypeptides contain signal sequences, indicating these polypeptides are capable of being expressed in one cell while exerting their effect in another cell. Accordingly, the NFIF polypeptides may be expressed by cells that then release NFIF polypeptide into the bloodstream or other transport system (lymph, etc.)
25 where they are transported to tissues where they exert their effect on NFB induction.

The discussion which follows describes how to identify compositions and methods for regulating NFB induction based on the present discovery of the NFIF polypeptides.

30

Compositions

The present invention provides compositions in a biologically compatible (biocompatible) solution comprising the polypeptides, nucleic acids, and vectors of the invention. A biologically compatible solution is a solution in which the polypeptide, nucleic acid or vector of the invention is maintained in an active form, e.g., in a form
5 able to effect a biological activity. For example, a polypeptide of the invention could have NFB activation or deactivation activity; an antibody (which is itself a polypeptide) would bind a polypeptide of the invention; a nucleic acid would be able to replicate, translate a message, or hybridize to a complementary nucleic acid; and a vector would be able to transfect a target cell. Generally, such a biologically compatible solution will
10 be an aqueous buffer, e.g., Tris, phosphate, or HEPES buffer, containing salt ions. Usually the concentration of salt ions will be similar to physiological levels. In a specific embodiment, the biocompatible solution is a pharmaceutically acceptable composition. Biologically compatible solutions may include stabilizing agents and preservatives.

15 Such compositions can be formulated for administration by topical, oral, parenteral, intranasal, subcutaneous, and intraocular, routes. Parenteral administration is meant to include intravenous injection, intramuscular injection, intraarterial injection or infusion techniques. The composition may be administered parenterally in dosage
20 unit formulations containing standard, well known nontoxic physiologically acceptable carriers, adjuvants and vehicles as desired.

The preferred sterile injectable preparations can be a solution or suspension in a nontoxic parenterally acceptable solvent or diluent. Examples of pharmaceutically
25 acceptable carriers are saline, buffered saline, isotonic saline (e.g. monosodium or disodium phosphate, sodium, potassium, calcium or magnesium chloride, or mixtures of such salts), Ringer's solution, dextrose, water, sterile water, glycerol, ethanol, and combinations thereof. 1,3-butanediol and sterile fixed oils are conveniently employed as solvents or suspending media. Any bland fixed oil can be employed including synthetic
30 mono- or di-glycerides. Fatty acids such as oleic acid also find use in the preparation of injectables.

The composition medium can also be a hydrogel which is prepared from any biocompatible or non-cytotoxic (homo or hetero) polymer, such as a hydrophilic polyacrylic acid polymer that can act as a drug absorbing sponge. Such polymers have been described, for example, in application WO93/08845, the entire contents of which
5 are hereby incorporated by reference. Certain of them, such as, in particular, those obtained from ethylene and/or propylene oxide are commercially available. A hydrogel can be deposited directly onto the surface of the tissue to be treated, for example, during surgical intervention.

10 Another preferred embodiment of the present invention relates to a pharmaceutical composition comprising a replication defective recombinant virus and poloxamer. More specifically, the invention relates to a composition comprising a replication defective recombinant virus comprising a nucleic acid encoding an NFIF polypeptide and poloxamer. A preferred poloxamer is Poloxamer 407, which is
15 commercially available (BASF, Parsippany, NJ) and is a non-toxic, biocompatible polyol, and is most preferred. A poloxamer impregnated with recombinant viruses may be deposited directly on the surface of the tissue to be treated, for example during a surgical intervention. Poloxamer possesses essentially the same advantages as a hydrogel while having a lower viscosity.

20 The present invention also relates to compositions useful in the manufacture of medicaments intended for the treatment of subjects affected with an NFB-regulated inflammatory response and compositions useful in the manufacture of medicaments intended to prevent subjects from being affected with an NFB-regulated inflammatory
25 response. Accordingly, the present invention includes the use of an NFIF polypeptide according to the invention for the manufacture of a medicament intended for the treatment and/or prevention of an NFB-regulated inflammatory response. The present invention also includes the use of a nucleic acid according to the invention, encoding an NFIF polypeptide, for the manufacture of a medicament intended for the treatment
30 and/or prevention of an NFB-regulated inflammatory response. The present invention further includes the use of a recombinant vector according to the invention, comprising a nucleic acid encoding an NFIF polypeptide, for the manufacture of a medicament

intended for the treatment and/or prevention of an NFB-regulated inflammatory response. The present invention also includes the use of a defective recombinant viral vector according to the invention for the manufacture of a medicament intended for the treatment and/or prevention of an NFB-regulated inflammatory response.

5

Another aspect of the present invention relates to the use of compositions comprising cells genetically modified *ex vivo* with a recombinant virus and compositions comprising cells containing such recombinant viruses which are implanted in the body, facilitating prolonged and effective expression *in vivo* of an NFIF polypeptide according to the invention.

10

Therapeutic Compounds Based on NFIF that Inhibit Activation of NFB

15

As explained above, in certain situations, it is desirable to reduce or inhibit NFB-regulated immune responses. By inhibiting the expression of the NFIF gene or interfering with the activity of NFIF polypeptides, it is possible to inhibit induction of NFB and consequently inhibit or prevent NFB-regulated immune responses that result in atherosclerosis and other diseases.

20

There are a variety of therapeutic compounds which may be used to inhibit NFIF expression or activity. These therapeutic compounds may be nucleic acids, polypeptides, peptides or non-peptide small molecules. In one embodiment, antisense nucleic acids are used to decrease expression of the NFIF gene by inhibiting processing (splicing) of the NFIF primary transcript. In another embodiment, ribozymes that cleave NFIF mRNA are used, preventing the synthesis of NFIF. Polypeptides of the present invention include antibodies or other binding proteins that bind NFIF polypeptides and interfere with their ability to activate NFB induction. In addition, compounds such as small molecule inhibitors which inhibit NFIF gene expression or the activity of NFIF polypeptides can be identified in reporter gene assays as described above and administered to patients to inhibit NFIF expression/activity. The same methods described above for introducing NFIF polypeptides and nucleic acids encoding NFIF

30

polypeptides into a patient's body are used to administer the various anti-NFIF compositions.

Antisense Nucleic Acids

5 The down regulation of gene expression using antisense nucleic acids can be achieved at the translational or transcriptional level. Antisense nucleic acids of the invention are preferably nucleic acid fragments capable of specifically hybridizing with all or part of a nucleic acid encoding NFIF or the corresponding messenger RNA. In addition, antisense nucleic acids may be designed or identified which decrease
10 expression of the NFIF gene by inhibiting splicing of its primary transcript. With knowledge of the structure and partial sequence of the NFIF gene, such antisense nucleic acids can be designed and tested for efficacy.

 The antisense nucleic acids are preferably oligonucleotides and may consist
15 entirely of deoxyribo-nucleotides, modified deoxyribonucleotides, or some combination of both. The antisense nucleic acids can be synthetic oligonucleotides. The oligonucleotides may be chemically modified, if desired, to improve stability and/or selectivity. Since oligonucleotides are susceptible to degradation by intracellular nucleases, the modifications can include, for example, the use of a sulfur group to
20 replace the free oxygen of the phosphodiester bond. This modification is called a phosphorothioate linkage. Phosphorothioate antisense oligonucleotides are water soluble, polyanionic, and resistant to endogenous nucleases. In addition, when a phosphorothioate antisense oligonucleotide hybridizes to its target site, the RNA-DNA duplex activates the endogenous enzyme ribonuclease (Rnase) H, which cleaves the
25 mRNA component of the hybrid molecule.

 In addition, antisense oligonucleotides with phosphoramidite and polyamide (peptide) linkages can be synthesized. These molecules should be very resistant to nuclease degradation. Furthermore, chemical groups can be added to the 2' carbon of
30 the sugar moiety and the 5 carbon (C-5) of pyrimidines to enhance stability and facilitate the binding of the antisense oligonucleotide to its target site. Modifications may include 2' deoxy, O-pentoxy, O-propoxy, O-methoxy, fluoro, methoxyethoxy phosphoro-

thioates, modified bases, as well as other modifications known to those of skill in the art.

The antisense nucleic acids can also be DNA sequences whose expression in the cell produces RNA complementary to all or part of the NFIF mRNA. Antisense nucleic acids can be prepared by expression of all or part of a sequence selected from the group consisting of the sequence in Figure 3, in the opposite orientation, as described in EP 140308. Any length of antisense sequence is suitable for practice of the invention so long as it is capable of down-regulating or blocking expression of NFIF. Preferably, the antisense sequence is at least 20 nucleotides in length. The preparation and use of antisense nucleic acids, DNA encoding antisense RNAs and the use of oligo and genetic antisense is disclosed in WO92/15680, the contents of which are incorporated herein by reference.

One approach to determining the optimum fragment of NFIF to use in an antisense nucleic acid treatment method involves preparing random fragments of NFIF cDNA by mechanical shearing, enzymatic treatment, and cloning the fragment into any of the vector systems described herein. Individual clones or pools of clones are used to infect NFIF-expressing cells, and effective antisense NFIF cDNA fragments are identified by monitoring NFIF expression at the RNA or protein level.

The retroviral, adeno-associated viral, and adenoviral vector systems discussed hereinabove may all be used to introduce and express antisense nucleic acids in cells. Antisense synthetic oligonucleotides may be introduced in a variety of ways, including the methods discussed hereinbelow.

Ribozymes

Reductions in the levels of NFIF polypeptide may be accomplished using ribozymes. Ribozymes are catalytic RNA molecules (RNA enzymes) that have separate catalytic and substrate binding domains. The substrate binding sequence combines by nucleotide complementarity and, possibly, nonhydrogen bond interactions with its target sequence. The catalytic portion cleaves the target RNA at a specific site. The substrate domain of a ribozyme can be engineered to direct it to a specified mRNA sequence. The

ribozyme recognizes and then binds a target mRNA through complementary base-pairing. Once it is bound to the correct target site, the ribozyme acts enzymatically to cut the target mRNA. Cleavage of the NFIF mRNA by a ribozyme destroys its ability to direct synthesis of NFIF polypeptide. Once the ribozyme has cleaved its target
 5 sequence, it is released and can repeatedly bind and cleave at other NFIF mRNAs.

Examples of ribozymes for use in the practice of the present invention include a variety of motifs, for example, hammerhead motif, hairpin motif, a hepatitis delta virus, group I intron or RnaseP RNA (in association with an RNA guide sequence) motif or
 10 Neurospora VS RNA motif. Hammerhead motifs are described by Rossi et al., 1992, *Aids Research and Human Retroviruses*, 8, 183. Hairpin motifs are described in Hampel and Tritz, 1989, *Biochemistry*, 28, 4929, and Hampel et al., 1990, *Nucleic Acids Res.*, 18, 299. The hepatitis delta virus motif is described by Perrotta and Been, 1992, *Biochemistry*, 31, 16, the RnaseP motif is described by Guerrier-Takada et al.,
 15 1983, *Cell*, 35, 849, the Neurospora VS RNA ribozyme motif is described by Collins (Saville and Collins, 1990, *Cell*, 61, 685-696; Saville and Collins, 1991, *Proc. Natl. Acad. Sci. USA*, 88, 8826-8830; Collins and Olive, 1993, *Biochemistry*, 32, 2795-2799) the Group I intron motif is described by Cech et al., U.S. Patent No. 4,987,071.

20 One approach in preparing a ribozyme is to synthesize chemically an oligodeoxyribonucleotide with a ribozyme catalytic domain (–20 nucleotides) flanked by sequences that hybridize to the target NFIF mRNA after transcription. The oligodeoxyribonucleotide is amplified by using the substrate binding sequences as primers. The amplification product is cloned into a eukaryotic expression vector.

25 Ribozymes possessing a hammerhead or hairpin structure are prepared readily since these catalytic RNA molecules can be expressed within cells from eukaryotic promoters (e.g., Scanlon et al., 1991, *Proc. Natl. Acad. Sci. USA*, 88, 10591-5; Kashani-Sabet et al., 1992, *Antisense Res. Dev.*, 2, 3-15; Dropulic et al., 1992, *J. Virol.*, 66, 1432-41; Weerasinghe et al., 1991, *J. Virol.*, 65, 5531-4; Ojwang et al.,
 30 1992, *Proc. Natl. Acad. Sci. USA*, 89, 10802-6; Chen et al., 1992, *Nucleic Acids Res.*, 20, 4581-9; Sarver et al., 1990, *Science*, 247, 1222-1225)). A ribozyme of the present

invention can be expressed in eukaryotic cells from the appropriate DNA vector. If desired, the activity of the ribozyme may be augmented by its release from the primary transcript by a second ribozyme (Ohkawa et al., 1992, *Nucleic Acids Symp. Ser.*, 27, 15-6; Taira et al., 1991, *Nucleic Acids Res.*, 19, 5125-30; Ventura et al., 1993, *Nucleic*
 5 *Acids Res.*, 21, 3249-55).

In one approach to preparing ribozymes, ribozymes are expressed from transcription units inserted into DNA, RNA, or viral vectors. Transcription of the ribozyme sequences are driven from a promoter for eukaryotic RNA polymerase I (pol
 10 (I), RNA polymerase II (pol II), or RNA polymerase III (pol III). Transcripts from pol II or pol III promoters will be expressed at high levels in all cells; the levels of a given pol II promoter in a given cell type will depend on nearby gene regulatory sequences. Prokaryotic RNA polymerase promoters are also used, providing that the prokaryotic RNA polymerase enzyme is expressed in the appropriate cells (Elroy-Stein and Moss,
 15 1990, *Proc. Natl. Acad. Sci. USA*, 87, 6743-7; Gao and Huang 1993, *Nucleic Acids Res.*, 21, 2867-72; Lieber et al., 1993, *Methods Enzymol.*, 217, 47-66; Zhou et al., 1990, *Mol. Cell. Biol.*, 10, 4529-37). It has been demonstrated that ribozymes expressed from these promoters can function in mammalian cells (Kashani-Sabet et al., 1992, *Antisense Res. Dev.*, 2, 3-15; Ojwang et al., 1992, *Proc. Natl. Acad. Sci. USA*,
 20 89, 10802-6; Chen et al., 1992 *Nucleic Acids Res.*, 20, 4581-9; Yu et al., 1993, *Proc. Natl. Acad. Sci. USA*, 90, 6340-4; L'Huillier et al., 1992, *EMBO J.*, 11, 4411-8; Lisiewicz et al., 1993, *Proc. Natl. Acad. Sci. USA*, 90, 8000-4).

In one embodiment of the present invention, a transcription unit expressing a
 25 ribozyme that cleaves NFIF RNA is inserted into a plasmid DNA vector, a retrovirus vector, an adenovirus DNA viral vector or an adeno-associated virus vector. The recombinant vectors are preferably DNA plasmids or adenovirus vectors. However, other mammalian cell vectors that direct the expression of RNA may be used for this purpose. The vectors are delivered as recombinant viral particles. DNA may be
 30 delivered alone or complexed with various vehicles. The DNA, DNA/vehicle complexes, or the recombinant virus particles are locally administered to the site of treatment, as discussed below. Preferably, recombinant vectors capable of expressing the

ribozymes are locally delivered as described below, and persist in target cells. Once expressed, the ribozymes cleave the target NFIF mRNA.

Ribozymes may be administered to a patient by a variety of methods. They may be added directly to target tissues, complexed with cationic lipids, packaged within liposomes, or delivered to target cells by other methods known in the art. Localized administration to the desired tissues may be done by catheter, infusion pump or stent, with or without incorporation of the ribozyme in biopolymers as discussed hereinbelow.

Alternative routes of delivery include, but are not limited to, intravenous injection, intramuscular injection, subcutaneous injection, aerosol inhalation, oral (tablet or pill form), topical, systemic, ocular, intraperitoneal and/or intrathecal delivery. More detailed descriptions of ribozyme delivery and administration are provided in Sullivan et al., PCT WO94/02595 and Draper et al., PCT WO93/23569, which are incorporated by reference herein.

15

Antibodies

The present invention provides antibodies against the NFIF polypeptide. These antibodies may be monoclonal antibodies or polyclonal antibodies. The present invention includes chimeric, single chain, and humanized antibodies, as well as Fab fragments and the products of an Fab expression library, and Fv fragments and the products of an Fv expression library.

Polyclonal antibodies may be prepared against an antigenic fragment of an NFIF polypeptide. Antibodies may also be generated against the intact NFIF protein or polypeptide, or against a fragment, derivative, or epitope of the protein or polypeptide. Antibodies may be obtained following the administration of the protein, polypeptide, fragment, derivative, or epitope to an animal, using the techniques and procedures known in the art.

Monoclonal antibodies may be prepared using the method of Mishell, B. B., et al., Selected Methods In Cellular Immunology, (W.H. Freeman, ed.) San Francisco (1980). Briefly, a polypeptide of the present invention is used to immunize spleen cells

of Balb/C mice. The immunized spleen cells are fused with myeloma cells. Fused cells containing spleen and myeloma cell characteristics are isolated by growth in HAT medium, a medium which kills both parental cells, but allows the fused products to survive and grow.

5

The monoclonal antibodies of the present invention may be "humanized" to prevent the host from mounting an immune response to the antibodies. A "humanized antibody" is one in which the complementarity determining regions (CDRs) and/or other portions of the light and/or heavy variable domain framework are derived from a non-
10 human immunoglobulin, but the remaining portions of the molecule are derived from one or more human immunoglobulins. Humanized antibodies also include antibodies characterized by a humanized heavy chain associated with a donor or acceptor unmodified light chain or a chimeric light chain, or vice versa. The humanization of antibodies may be accomplished by methods known in the art (see, e.g. G.E. Mark and
15 E.A. Padlan, "Chapter 4. Humanization of Monoclonal Antibodies", The Handbook of Experimental Pharmacology Vol. 113, Springer-Verlag, New York, 1994). Transgenic animals may be used to express humanized antibodies.

Techniques known in the art for the production of single chain antibodies can be
20 adapted to produce single chain antibodies to the immunogenic polypeptides and proteins of the present invention.

In a preferred embodiment, an anti-NFIF antibody is used to bind to and inhibit the activity of NFIF in a patient.

25

The anti-NFIF antibodies are also useful in assays for detecting or quantitating levels of NFIF. In one embodiment, these assays provide a clinical diagnosis and assessment of NFIF in various disease states and a method for monitoring treatment efficacy. An example of an anti-NFIF antibody used to bind NFIF and identify its
30 presence in tissues is provided in Example 5. These anti-NFIF antibodies may additionally be used to quantitate NFIF in a tissue sample.

Methods of Treatment

The present invention provides methods of treatment which comprise the administration to a human or other animal of an effective amount of a composition of the invention.

5

Effective amounts may vary, depending on the age, type and severity of the condition to be treated, body weight, desired duration of treatment, method of administration, and other parameters. Effective amounts are determined by a physician or other qualified medical professional. In most cases, the dosage levels may be adjusted so that the desired levels of NFIF or other therapeutic compounds can be achieved and maintained.

10

Polypeptides according to the invention are generally administered in doses of about 0.01 mg/kg to about 100 mg/kg, more preferably about 0.1 mg/kg to about 50 mg/kg, and even more preferably about 1 mg/kg to about 10 mg/kg of body weight per day.

15

Neutralizing antibodies according to the invention may be delivered as a bolus only, infused over time or both administered as a bolus and infused over time. Although the dosage amount will vary based on the parameters above, and on the binding ability of the antibody, a dose 0.2 to 0.6 mg/kg may be given as a bolus followed by a 2 to 12 hour infusion period. Alternatively, multiple bolus injections are administered every other day or every third or fourth day as needed. Dosage levels may be adjusted as determined by NFIF levels and/or NFB induction levels.

20

25

As discussed hereinabove, recombinant viruses may be used to introduce both DNA encoding NFIF and subfragments of NFIF as well as antisense nucleic acids. Recombinant viruses according to the invention are generally formulated and administered in the form of doses of between about 10^4 and about 10^{14} pfu. In the case of AAVs and adenoviruses, doses of from about 10^6 to about 10^{11} pfu are preferably used. The term pfu ("plaque-forming unit") corresponds to the infective power of a suspension of virions and is determined by infecting an appropriate cell culture and

30

measuring the number of plaques formed. The techniques for determining the pfu titre of a viral solution are well documented in the prior art.

Ribozymes according to the present invention may be administered in amounts
5 ranging from about 5 to about 50 mg/kg/day in a pharmaceutically acceptable carrier. Dosage levels may be adjusted based on the measured therapeutic efficacy.

Appropriate levels of inhibitor or enhancer molecules may be determined by qualified medical personnel using the parameters discussed above.

10

Methods for Increasing the Level of NFIF Polypeptide Activity

The methods for increasing the expression or activity of NFIF polypeptide include, but are not limited to, administration of a composition comprising the NFIF polypeptide, administration of a composition comprising an expression vector which
15 encodes the NFIF polypeptide, administration of a composition comprising an enhancer molecule which enhances the activity of the NFIF polypeptide and administration of an enhancer molecule which increases expression of the NFIF gene.

Methods Utilizing NFIF Polypeptides

20 In one embodiment, the level of NFIF activity is increased through the administration of a composition comprising the NFIF polypeptide. This composition may be administered in a convenient manner, such as by the oral, topical, intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal, or intradermal routes. The composition may be administered directly or it may be encapsulated (e.g. in a lipid
25 system, in amino acid microspheres, or in globular dendrimers). The polypeptide may, in some cases, be attached to another polymer such as serum albumin or polyvinyl pyrrolidone.

Methods Utilizing Vectors that Express NFIF

30 In another embodiment, the level of NFIF is increased through the use of gene therapy, that is, through the administration of composition comprising a nucleic acid which encodes and directs the expression of the NFIF polypeptide. In this embodiment,

the NFIF polypeptide is cloned into an appropriate expression vector. Possible vector systems and promoters are extensively discussed above. The expression vector is transferred into the target tissue using one of the vector delivery systems discussed above. This transfer is carried out either *ex vivo* in a procedure in which the nucleic acid is transferred to cells in the laboratory and the modified cells are then administered to the human or other animal, or *in vivo* in a procedure in which the nucleic acid is transferred directly to cells within the human or other animal. In preferred embodiments, an adenoviral vector system is used to deliver the expression vector. If desired, a tissue specific promoter is utilized in the expression vector as described above.

Non-viral vectors may be transferred into cells using any of the methods known in the art, including calcium phosphate coprecipitation, lipofection (synthetic anionic and cationic liposomes), receptor-mediated gene delivery, naked DNA injection, electroporation and bioballistic or particle acceleration.

Methods Utilizing an Enhancer Molecule which Enhances the Activity of NFIF

In another embodiment, the activity of NFIF is enhanced by enhancer molecules that increase the activity of NFIF or increase its appropriate recognition by cellular binding sites. These enhancer molecules may be introduced by the same methods discussed above for the administration of polypeptides.

Methods Utilizing an Enhancer Molecule which Increases NFIF Gene Expression

In another embodiment, the level of NFIF is increased through the use of small molecular weight compounds, which can upregulate NFIF expression at the level of transcription, translation, or post-translation. These compounds may be administered by the same methods discussed above for the administration of polypeptides.

30

Methods for Treating or Preventing an NFB-regulated Inflammatory Response

The present invention includes methods for the treatment or prevention of NFB-regulated inflammatory responses including, but not limited to, rheumatoid arthritis, atherosclerosis, autoimmune diseases, viral diseases, NSAID-induced gastropathy, neurodegenerative diseases, scrapie, sepsis, apoptosis, Crohn=s disease, renal disease, restenosis, brain injury/inflammation, Alzheimer=s disease, asthma, and improperly regulated expression of pleiotropic cytokines.

These methods include, but are not limited to, administration of a composition comprising an antisense nucleic acid, administration of a composition comprising an intracellular binding protein such as an antibody, administration of an inhibitory molecule which inhibits the activity of NFIF, for example, a composition comprising an expression vector encoding a subfragment of NFIF or a small molecular weight molecule, including administration of a small molecular weight compound which down regulates NFIF expression at the level of transcription, translation or post-translation, administration of a ribozyme which cleaves mRNA encoding NFIF, administration of a medicament manufactured using an NFIF polypeptide, administration of a medicament manufactured using a nucleic acid encoding an NFIF polypeptide, administration of a medicament manufactured using a recombinant vector which includes nucleic acid encoding an NFIF polypeptide and administration of a medicament manufactured using a defective recombinant viral vector which includes nucleic acid encoding an NFIF polypeptide.

Methods for Lowering Levels of NFIF Polypeptide Activity

The methods for decreasing the expression of NFIF polypeptide in order to decrease NFB induction include, but are not limited to, administration of a composition comprising an antisense nucleic acid, administration of a composition comprising an intracellular binding protein such as an antibody, administration of an inhibitory molecule which inhibits the activity of NFIF, for example, a composition comprising an expression vector encoding a subfragment of NFIF or a small molecular weight molecule, including administration of a small molecular weight compound which down

regulates NFIF expression at the level of transcription, translation or post-translation, and administration of a ribozyme which cleaves mRNA encoding NFIF.

Methods Utilizing Antisense Nucleic Acids

5 In one embodiment, a composition comprising an antisense nucleic acid is used to down-regulate or block the expression of NFIF. In one preferred embodiment, the nucleic acid encodes antisense RNA molecules. In this embodiment, the nucleic acid is operably linked to signals enabling expression of the nucleic acid sequence and is introduced into a cell utilizing, preferably, recombinant vector constructs, which will
10 express the antisense nucleic acid once the vector is introduced into the cell. Examples of suitable vectors includes plasmids, adenoviruses, adeno-associated viruses, retroviruses, and herpes viruses. Preferably, the vector is an adenovirus. Most preferably, the vector is a replication defective adenovirus comprising a deletion in the E1 and/or E3 regions of the virus.

15 In another embodiment, the antisense nucleic acid is synthesized and may be chemically modified to resist degradation by intracellular nucleases, as discussed above. Synthetic antisense oligonucleotides can be introduced to a cell using liposomes. Cellular uptake occurs when an antisense oligonucleotide is encapsulated within a
20 liposome. With an effective delivery system, low, non-toxic concentrations of the antisense molecule can be used to inhibit translation of the target mRNA. Moreover, liposomes that are conjugated with cell-specific binding sites direct an antisense oligonucleotide to a particular tissue.

Methods Utilizing Neutralizing Antibodies and Other Binding Proteins

25 In another embodiment, the expression of NFIF is down-regulated or blocked by the expression of a nucleic acid sequence encoding an intracellular binding protein which is capable of selectively interacting with NFIF. WO 94/29446 and WO 94/02610, the contents of which are incorporated herein by reference, disclose cellular transfection
30 with genes encoding an intracellular binding protein. An intracellular binding protein includes any protein capable of selectively interacting, or binding, with NFIF in the cell in which it is expressed and of neutralizing the function of bound NFIF. Preferably, the

intracellular binding protein is a neutralizing antibody or a fragment of a neutralizing antibody. More preferably, the intracellular binding protein is a single chain antibody.

WO 94/02610 discloses preparation of antibodies and identification of the nucleic acid encoding a particular antibody. Using NFIF or a fragment thereof, a specific monoclonal antibody is prepared by techniques known to those skilled in the art. A vector comprising the nucleic acid encoding an intracellular binding protein, or a portion thereof, and capable of expression in a host cell is subsequently prepared for use in the method of this invention.

Alternatively, NFIF activity can be blocked by administration of a neutralizing antibody into the circulation. Such a neutralizing antibody can be administered directly as a protein, or it can be expressed from a vector (with a secretory signal).

Methods Utilizing an Inhibitory Molecule which Prevents NFIF Gene Expression

In another embodiment, inhibitory molecules, including small molecular weight compounds, are able to down regulate NFIF expression at the level of transcription, translation or post-translation. In order to identify such inhibitory molecules, the reporter gene systems described above may be used. These inhibitory molecules may be combined with a pharmaceutically acceptable carrier and administered using conventional methods known in the art.

Methods Utilizing Ribozymes

Ribozymes may be administered to cells by encapsulation in liposomes, by iontophoresis, by incorporation into hydrogels, cyclodextrins, biodegradable nanocapsules, and bioadhesive microspheres or by any of a variety of other methods discussed above. The ribozyme may be delivered to a target tissue by direct injection or by use of a catheter, infusion pump or stent. Alternative routes of delivery include intravenous injection, intramuscular injection, subcutaneous injection, aerosol inhalation, oral (tablet or pill form), topical, systemic, ocular, intraperitoneal and/or intrathecal delivery.

In preferred embodiments, a ribozyme-encoding sequence is cloned into a DNA expression vector. Transcription of the ribozyme sequence is driven from an eukaryotic RNA polymerase II (pol II), or RNA polymerase III (pol III) promoter. The expression
5 vector can be incorporated into a variety of vectors including the viral DNA vectors such as adenovirus or adeno-associated virus vectors discussed above.

In a preferred embodiment of the invention, a transcription unit expressing a ribozyme that cleaves NFIF RNA is inserted into an adenovirus DNA viral vector. The
10 vector is delivered as recombinant viral particles and is locally administered to the site of treatment, through the use of a catheter, stent or infusion pump.

Examples

Example 1 - Method for Cloning NFIF-14b and NFIF-7a

15 To isolate the DNA encoding NFIF-14b and NFIF-7a, a cDNA library was prepared from human vascular smooth muscle cell (SMC) total RNA using a Clontech MarathonJ cDNA library synthesis kit. The cDNA synthesis kit also includes a control human placenta DNA library. The libraries are linear and amplifiable by PCR.

20 The NFIF-14b and NFIF-7a genes were obtained by PCR performed on both Human Vascular SMC and Human Placenta cDNA marathon libraries.

To perform the PCR reactions, standard PCR components [Perkin Elmer 10X buffer and Taq Gold Polymerase] were used and the reaction was performed as follows.
25 The reaction mixtures were heated to 94EC for 10 minutes, then thermocycled 10 times with a denaturing step of 94EC for 30 seconds, an annealing step of 65EC for 30 seconds, and an extension step of 72EC for 1.5 minutes. Following these ten cycles, the reaction was thermocycled 30 times with a denaturing step of 94EC for 30 seconds, an annealing step of 46EC for 30 seconds., and an extension step of 72EC for 1.5 minutes.
30 Following these 30 cycles, the reactions were incubated at 72EC for 7 minutes and stored at 4EC. The PCR reactions were performed in a Perkin Elmer 9600 thermocycler.

The PCR product from each PCR was ligated into pCR2.1 (Invitrogen) and transformed in DH5-E.coli. Clones were picked and grown up in 5 ml of LB-amp; plasmids were then isolated and digested with EcoRI to check for the size of the released insert. Five out of fifteen clones gave bands of size 1.2-1.3 Kb, suggestive of the target gene and were sequenced.

Putative NFIF clones were identified based on an open reading frame and sequence alignment of the sequence against a predicted 5' sequence. Clones Nos. 14b and 7a were chosen for further characterization.

Large plasmid preparations were grown up from original glycerol stocks of clone 7 and 14, and isolated using a Qiagen MaxiPrep Kit.

Example 2 - Subcloning of clone 7a and 14b into a eukaryotic expression vector

Clones 14b and 7a were further subcloned into the plasmid vector pcDNA3.1myc-his available from Invitrogen. This expression vector includes a strong promoter for high level expression in mammalian cells and a selection marker for generating stable cell lines. A C-terminal fusion tag in the vector features a polyhistidine sequence for rapid purification and a myc epitope for convenient detection with an anti-myc antibody.

PCR reactions were performed using PCR primers

5'hasm - 5'-tccaccatggcgctggtgcgcgcactc-3' and

fushasm3' - 3'-ctggatatcgtaattgtgctttatataaagctg-5' and the pCR2.1 construct for each full length clone as template. PCR conditions using 475pg of pCR2.1 clone 7a or 500 pg of pCR2.1 clone 14b were as follows: the reaction mixtures were heated to 95EC for 10 minutes, then thermocycled 30 times with a denaturing step of 95EC for 30 seconds, an annealing step of 52EC for 30 seconds, and then an extension step of 72EC for 1.5 minutes. Following these 30 cycles, the reactions were incubated at 72EC for 7 minutes, and stored at 4EC.

The PCR product was ligated into pCR2.1 and the insert DNA sequenced utilizing a Big Dye Terminator Cycle Sequencing Ready Reaction and a Perkin Elmer Applied Biosystems ABI Prism 377 DNA Sequencer. Sequencing demonstrated that the PCR primer removed the natural stop codon (TAG) and introduced an EcoRV site.

5

The positive constructs of NFIF-14b and -7a were digested with EcoRI and EcoRV and subcloned into pcDNA3.1 mychis (Invitrogen) cut with EcoRI and EcoRV. This resulted in the myc tag being fused to the 3' end of the NFIF cDNAs. These constructs were called pcDNA3.1mychasm7a and pcDNA3.1hasm14bmyc.

10

Example 3 - Preparation of Deletion Variants of NFIF

Deletion mutants of NFIF-14b and NFIF-7a were prepared by PCR reactions using the pcDNA3.1mychasm7a and pcDNA3.1hasm14bmyc plasmids described above.

Two sets of PCR primers were used to prepare the deletion mutants:

5 hasm313mut + hasm3'mut:
 5' gctccaccatgatatggacaggggatag 3'
 5' gccactgtgctggatcgtataataac 3'
 hasm396mut + hasm3'mut:
 5' gctccaccatgacaaccaccatccagagtc 3'
 10 5' gccactgtgctggatcgtataataac 3'

The hasm313 + hasm3'mut primer pair produced the identical sequence of the full length clone 14b or 7a cDNAs but on an open reading frame initiated downstream at bp313 (ATG). (Numbering is from the ATG in the full length clone.) The consensus
 15 Kozak sequence was included in the forward primer to optimize translation.

The hasm396 + hasm3'mut primer pair produced the identical sequence of the full length cDNAs but on an open reading frame initiated downstream at bp 394 (ATG).

20 The PCR products were ligated into pCR2.1 and the insert DNA sequenced. Each deletion mutant was cut with EcoRI/EcoRV and subcloned into pcDNA3.1mychis cut with EcoRI and EcoRV. NFIF-14b deletion clones with the correct sequence were named pcDNAMychis 14-313 and pcDNAMychis 14-396. NFIF-7a deletion clones with the correct sequence were named pcDNAMychis 7-313, and pcDNAMychis 7-396.

25 Analysis of the above clones using in vitro transcription/translation incorporating [³⁵S methionine] into the reaction [Promega TnT Quick Coupled Transcription Translation kit # L1170] produced bands of the size expected(NFIF 14b=51 kDa; NFIF 14-313=40 kDa; NFIF 14-396=37 kDa; NFIF 7a=41 kDa; NFIF 7-313=30 kDa; and
 30 NFIF 7-396=27 kDa) for shortened open reading frames. The facility of translation was important to determine before assessing functional activity and determining which domains were important for activity.

**Example 4 - Method for transfecting cells with NFIF-14b
or NFIF-7a to produce stable cell lines containing plasmids**

5 To prepare stable cell lines containing NFIF-14b, NFIF-7a and the deletion mutants, Falcon 6 well plates were seeded with 2×10^5 HEK293 or COS-7 cells, the finalized cDNA NFIF constructs pcDNAmychis7, pcDNAmychis7-313, pcDNAmychis7-396; pcDNAmychis14, pcDNAmychis14-313, pcDNAmychis14-396 (0.8 ug) along with the Stratagene pNFB-Luc reporter gene vector (0.1 ug) and Clontech
10 EGFP vector (0.1 ug), were transfected using 6 ul Lipofectamine and 200 ul Optimem into the 6 well plates (with 800 ul fresh Optimem/well). The transfection was incubated for 4 hr at 37EC and then the plates were fed with complete media (3ml).

At 24, 48 and 72 hours post transfection lysates were made using 200 ul 1X
15 Reporter Lysis Buffer solution (E397A-Promega). Lysates were incubated for 20 minutes on ice, vortexed, then centrifuged at 12,000 rpm for 5 minutes. The supernatant was used to assess luciferase activity per the instructions in the Stratagene pNFB-luc reporter gene kit.

20 To prepare stable cells lines, duplicates of the above 48 hr cultures were split at a 1:100 ratio into complete growth media along with G418 for selection of cells incorporating the plasmids in their DNA. Single cell clones were isolated and transferred into 48 well culture dishes for growout. Clones were chosen according to their activity in the NFB luciferase reporter assay.

25 Luciferase activity was measured from the cell lysates of all the transient infected cell lines. 20 l of lysate was added to a 96 well Costar serocluster (#3789) white round bottom plate. 100 l of Luciferase Reagent (Promega E1501) was added directly before reading the sample in a 1450 Microbeta Walko jet.

30 Analysis of the results from 24 hour, 48 hour, 72 hour timepoints identified the 48 hour timepoint as the most optimal for assessing activity. The positive control for the

assay was a 24 hour tumor necrosis factor (TNF) stimulation of the cells one day posttransfection with the pcDNA3.1mychis vector. In HEK293 cells, the full-length pcDNAmychis 7, pcDNAmychis14 cDNAs transient transfections demonstrated luminometer count per second at -3.1 fold and -2.1 fold respectively, above vector only (Figure 5) and in COS-7 cells (Figure 6) full-length clone pcDNAmychis 7 gave a 2.4 fold signal and pcDNAmychis 14 a 2.3 fold increase in luminometer count per second above vector only.

The deletion mutants (pcDNAmychis 7-313, pcDNAmychis 7-396, pcDNAmychis 14-313 and pcDNAmychis 14-396) showed reduced activity. Assays run on other days using the 48 hour timepoint gave variability in response ranging between a 2-5 fold increase for clone 7 (full-length) and 2-4 fold for clone 14.

Example 5 - Identification of Tissues Expressing or Containing NFIF Proteins

In order to determine which human tissues express the NFIF proteins, a Clontech pre-made Northern Blot (Human 12 lane #7780-1) was probed with randomly primed p32 labelled probes prepared from NFIF. The results can be seen in Figure 8. As Figure 8 illustrates, expression of NFIF is particularly evident in skeletal muscle, kidney, liver and placental tissue.

In order to determine if the NFIF protein was associated with pathologies including atherosclerosis that involve inflammation and to identify tissues that may be treated using the methods of the present invention, an immunocytochemical study was performed using a rabbit monoclonal antibody designated 99-06 directed against a peptide antigen (SKGANASNPFGPDV) derived from residues 65 to 79 of the NFIF protein. The peptide was synthesized at the 0.25 mmole scale using a solid phase methodology FMOC (9-fluorenylmethyloxycarbonyl) protection scheme in conjunction with the HOBt/HBTU activation chemistry (Fields et al., *Peptide Research*, 4:95-101 (1991)). An Applied Biosystems 433 Peptide Synthesizer running Applied Biosystems Fast-Moc coupling cycle was used for the synthesis of the peptide.

The peptide was cleaved for 1.5 hours at room temperature using a cleavage reagent of 82.5 % trifluoroacetic acid (TFA), 5 % phenol, 5 % H₂O, 5 % thioanisole, and 2.5 % ethanedithiol (King et al., *Intl. J. Peptide and Protein Research*, 36, 255-266 (1990)). Following cleavage, the peptide was precipitated with tert-butyl ether, washed, then dried for 1 hour under vacuum. The peptide was then solubilized in 0.1 % TFA/water and purified using C18 reverse phase HPLC. A purity level of >95 % was achieved for the peptide, along with correct MALDI-TOF molecular weight data.

10-30 mg of purified peptide was conjugated to keyhole limpet hemocyanin using 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide (EDC). The preparation was dialyzed against PBS at 4EC exhaustively. The hapten/carrier preparation was mixed 1:1 with Freund's Complete adjuvant and used to inject rabbits for the production of antisera against the peptide. The 15-residue peptide was used to prepare anti-NFIF antibodies in rabbits using standard methods known in the art. Rabbits were immunized by subcutaneous injection with an emulsion of antigen/complete Freund's adjuvant (50 % antigen solution/50 % complete Freund's adjuvant). Complete Freund's adjuvant (CFA) was used for the initial immunization because this adjuvant has been used very successfully in producing antibody responses to a diverse range of protein and peptide antigens and the incidence of adverse reactions is minimal to none. The injection schedule described below has been optimized for use with Freund's adjuvant. The area to be injected was shaved in order to permit visual assessment of any reaction to the emulsion. Prior to injection, the injection sites were cleansed with diluted Nolvasan scrub solution. Injection sites were on the back starting between the shoulder blades. The injection sites were widely separated and kept close to the midline to reduce the possibility of the animal scratching at the injection sites. Typically, 100 μ l containing 20 g of antigen were injected in 5 subcutaneous sites per rabbit for a total of 100 g/rabbit. Subsequent, booster injections were given at 4 week intervals with an emulsion of antigen/incomplete Freund's adjuvant. After six weeks, 5-30 mls of blood was collected to determine the titer of the antibody. If the antibody titer was not sufficient, additional booster injections were given at week intervals and blood samples collected every 2-4 weeks. Prior to each antigen injection, the condition of the previous injection

sites were evaluated for adverse reactions. Animals giving a positive antibody response were maintained and boosted and bled periodically in order to generate a large stock of antisera.

5 Antibody titration experiments were conducted with Antibody 99-06 to establish concentrations that yield minimal background and maximum detection of signal. A serial dilution study demonstrated the highest signal-to-noise ratio at a dilution of 1:500 and 1:1,000, and slides incubated with these concentrations of antibody were analyzed by the pathologists. Antibody 99-06 was used as the primary antibody, and the principal
10 detection system consisted of a Vector ABC-AP Kit (AK5002) with a Vector Red substrate kit that produced a fuschia-colored red deposit (SK-5100). Tissues were also stained with a positive control antibody (CD31) directed against the human leukocyte antigen (Stockinger et al., *J. Immunol.*, 145(11):3889-97 (1990)) to ensure that the tissue antigens were preserved and accessible for immunocytochemical analysis. In addition to
15 coronary artery samples, additional tissues that were stained and imaged during Phase II of this study were normal spleen, thymus, and lymph node. Staining was performed in the spleen, thymus and lymph node in order to assist in the identification of inflammatory subsets of cells that expressed this protein.

20 Within the spleen, thymus, and lymph node, Antibody 99-06 showed strong positive staining within the lymphocytes of the periarterial lymphatic sheath, the cords of Billroth, cortical thymic lymphocytes, germinal center cells in the lymph node, and within vascular endothelium and smooth muscle. Other cell types that stained moderately to strongly positive with this antibody included Schwann cells and cardiac
25 myocytes. The signal was predominantly nuclear in the cell types examined, except for cytoplasmic granularity that was seen within smooth muscle cells, and in macrophages that were filled with cellular debris in regions of plaque or organizing thrombus.

30 Within normal coronary arteries, the antibody was largely negative in the endothelium of the tunica intima except for occasional cells showing a nuclear signal, and myointimal smooth muscle cells showed faint nuclear signals. Within the tunica media, the antibody was moderately uniformly positive within smooth muscle cells. In

the adventitia, the endothelium, vascular smooth muscle and inflammatory cells were moderately positive for staining. The internal and external elastic lamellae were negative for staining.

5 Within the minimal atherosclerosis samples, the endothelium showed only occasional faint signals even in that overlying the atheroma. Within the superficial atheroma, the myointimal smooth muscle cells showed only faint signals. In deeper regions of the atheroma, proliferating fibroblastic forms and foamy macrophages showed an increased level of staining. In some samples that showed plaques containing
10 cholesterol and areas of calcification, the smooth muscle cells and foamy histiocytes surrounding the cholesterol-laden plaques and calcified material showed increased levels of staining.

 In the samples of moderate atherosclerosis, the endothelium, both in the tunica
15 intima and in areas of neovascularization, showed increased staining with Antibody 99-06. Inflammatory cells marginating along or subjacent to the endothelium were strongly positive for staining. Endothelium lining new vessels was moderately to strongly positive for staining, and in areas, the subendothelial myofibroblasts were less positive than associated endothelium. In general, fibroblastic forms of myointimal cells were
20 much less positive than either proliferating or histiocytic forms of smooth muscle cells. In addition, macrophages associated with cholesterol, calcification, or foamy macrophages within plaques showed an increased level of staining. The smooth muscle of the tunica media was fairly uniformly moderately positive within all samples in which the media was intact, and smooth muscle within adventitial vessels was similarly
25 positive.

 In cases of severe atherosclerosis, much of the endothelium lining the tunica intima was either denuded or the residual endothelium was faintly to moderately positive for staining. The plaque-associated inflammatory infiltrate showed very strong positive
30 staining within macrophages and lymphocytes, including within regions of neovascularization. Increased staining was also seen within macrophages and lymphocytes adjacent to or surrounding calcifications and cholesterol and within areas of

organizing thrombus. The endothelium lining new vessels was strongly positive in these areas. Increased signal intensity was again seen within the histiocytic and proliferating forms compared to fibroblastic forms of smooth muscle cells. In these cases, inflammatory cells such as a subset of lymphocytes, plasma cells, and macrophages within the adventitia were also strongly positive for staining.

In late stage areas of severe atherosclerosis with minimal inflammation and quiescent plaques, there was much less staining. In these samples, the endothelium was less intensely positive for staining, although residual lymphocytes and macrophages surrounding calcified material showed positive staining.

Biological Material Useful in Practicing the Invention

A deposit of Human NFB inducing factor 7a (NFIF-7a) cloned from a human vascular smooth muscle cell cDNA library subcloned into a pCR2.1 plasmid vector and a deposit of Human NFB inducing factor 14b (NFIF-14b) cloned from a human vascular smooth muscle cell cDNA library subcloned into a pCR2.1 plasmid vector will be made with the American Type Culture Collection. Access to these cultures will be available during pendency of the patent application to one determined by the Commissioner to be entitled thereto under 35 U.S.C. ' 122 and 37 C.F.R. ' 1.14.

We claim:

1. A nucleic acid encoding NFIF-14b polypeptide comprising an amino acid sequence as shown in Figure 1 (SEQ ID NO: 1).

5

2. A nucleic acid encoding NFIF-7a polypeptide comprising an amino acid sequence as shown in Figure 2 (SEQ ID NO: 2).

3. The nucleic acid of Claim 1 wherein said DNA is a cDNA.

10

4. The nucleic acid of Claim 2 wherein said DNA is a cDNA.

5. An isolated and purified NFIF-14b polypeptide which induces NFB and comprising an amino acid sequence as shown in Figure 1 (SEQ ID NO: 1).

15

6. An isolated and purified NFIF-7a polypeptide which induces NFB and comprising an amino acid sequence as shown in Figure 2 (SEQ ID NO: 2).

7. A method of increasing expression of NFB in a patient comprising
20 introducing into the body of said patient a composition that induces NFB.

8. The method of Claim 7 wherein said composition comprises an expression vector comprising a nucleic acid encoding NFIF-14b polypeptide.

25 9. The vector of Claim 8 selected from the group consisting of retroviral vectors, adenoviral vectors, adeno-associated viral vectors, herpesviral vectors, and naked DNA vectors.

30 10. The method of Claim 7 wherein said composition comprises an expression vector comprising a nucleic acid encoding NFIF-7a polypeptide.

11. The vector of Claim 10 selected from the group consisting of retroviral vectors, adenoviral vectors, adeno-associated viral vectors, herpesviral vectors, and naked DNA vectors.

5 12. The method of Claim 7 wherein said composition comprises a NFIF-14b polypeptide and a pharmaceutically acceptable carrier.

13. The method of Claim 7 wherein said composition comprises a NFIF-7a polypeptide and a pharmaceutically acceptable carrier.

10 14. A composition for lowering the expression of the NFIF gene in a patient comprising an antisense nucleic acid.

15 15. A composition for lowering the activity of an NFIF polypeptide in a patient comprising a neutralizing antibody that binds to an NFIF polypeptide and lowers its activity.

16. A composition for lowering the expression of NFIF in a patient comprising a ribozyme that cuts RNA encoding an NFIF polypeptide.

20 17. A method for evaluating whether a test compound is effective in inhibiting the activity of NFIF-14b based on the expression of an NFB-regulated reporter gene comprising:

25 (A) comparing the level of NFB-regulated gene expression in a first sample comprising: (1) NFIF-14b; (2) said NFB-regulated reporter gene; and (3) said test compound with the level of gene expression in a second sample comprising (4) NFIF-14b; and (5) said NFB-regulated reporter gene; and

(B) determining whether the expression of said reporter gene is lower in said first sample relative to said second sample.

18. A method for evaluating whether a test compound is effective in inhibiting the activity of NFIF-7a based on expression of an NFB-regulated reporter gene comprising:

(A) comparing the level of NFB-regulated gene expression in a first sample comprising: (1) NFIF-7a; (2) said NFB-regulated reporter gene; and (3) said test compound with the level of gene expression in a second sample comprising: (4) NFIF-7a; and (5) said NFB-regulated reporter gene; and

(B) determining whether the expression of said reporter gene is lower in said first sample relative to said second sample.

10

19. A method for identifying whether a test compound can enhance the activity of NFIF-14b based on the expression of an NFB-regulated reporter gene comprising:

(A) comparing the level of NFB-regulated gene expression in a first sample comprising (1) NFIF-14b; (2) said NFB-regulated reporter gene; and (3) said test compound with the level of gene expression in a second sample comprising: (4) NFIF-14b; and (5) said NFB-regulated reporter gene; and

(B) determining whether the expression of said reporter gene is higher in said first sample relative to said second sample.

20

20. A method for identifying compounds which enhance the activity of NFIF-7a based on the expression of an NFB-regulated reporter gene comprising:

(A) comparing the level of NFB-regulated gene expression in a first sample comprising: (1) NFIF-7a; (2) said NFB-regulated reporter gene; and (3) said test compound with the level of gene expression in a second sample comprising: (4) NFIF-7a; and (5) said NFB-regulated reporter gene; and

(B) determining whether the expression of said reporter gene is higher in said first sample relative to said second sample.

30

21. A method of inhibiting expression of NFB-dependent genes comprising administration to a patient of a composition that inhibits the activity of NFIF-14b.

22. A method of inhibiting expression of NFB-dependent genes comprising administration to a patient of a composition that inhibits the activity of NFIF-7a.

23. A method of inhibiting inflammation comprising administration of a composition that inhibits the activity of NFIF-14b.

24. A method of inhibiting inflammation comprising administration of a composition that inhibits the activity of NFIF-7a.

25. Use of an isolated NFIF polypeptide comprising an amino acid sequence of Figure 1 (SEQ ID NO: 1) for the manufacture of a medicament intended for the treatment and/or prevention of an NFB-regulated inflammatory response.

26. Use of a nucleic acid encoding an NFIF polypeptide comprising an amino acid sequence of Figure 1 (SEQ ID NO: 1) for the manufacture of a medicament intended for the treatment and/or prevention of an NFB-regulated inflammatory response.

27. Use of a recombinant vector comprising a nucleic acid encoding an NFIF polypeptide comprising an amino acid sequence of Figure 1 (SEQ ID NO: 1) for the manufacture of a medicament intended for the treatment and/or prevention of an NFB-regulated inflammatory response.

28. Use of a defective recombinant viral vector comprising a nucleic acid encoding an NFIF polypeptide comprising an amino acid sequence of Figure 1 (SEQ ID NO: 1) for the manufacture of a medicament intended for the treatment and/or prevention of an NFB-regulated inflammatory response.

29. Use of an isolated NFIF polypeptide comprising an amino acid sequence of Figure 2 (SEQ ID NO: 2) for the manufacture of a medicament intended for the treatment and/or prevention of an NFB-regulated inflammatory response.

30. Use of a nucleic acid encoding an NFIF polypeptide comprising an amino acid sequence of Figure 2 (SEQ ID NO: 2) for the manufacture of a medicament intended for the treatment and/or prevention of an NFB-regulated inflammatory response.

5

31. Use of a recombinant vector comprising a nucleic acid encoding an NFIF polypeptide comprising an amino acid sequence of Figure 2 (SEQ ID NO: 2) for the manufacture of a medicament intended for the treatment and/or prevention of an NFB-regulated inflammatory response.

10

32. Use of a defective recombinant viral vector comprising a nucleic acid encoding an NFIF polypeptide comprising an amino acid sequence of Figure 2 (SEQ ID NO: 2) for the manufacture of a medicament intended for the treatment and/or prevention of an NFB-regulated inflammatory response.

15

Abstract

The present invention is directed to nuclear factor B (NFB)-inducing factor polypeptides (NFIF polypeptides) which are capable of inducing NFB. The present invention includes within its scope NFIF polypeptides, including NFIF-14b and NFIF-7a, DNA, including cDNA, encoding these polypeptides, and expression vectors capable of expressing NFIF polypeptides. Also included are methods and compositions for increasing NFB induction in a patient, methods and compositions for lowering NFB induction in a patient, methods for inhibiting inflammation, and methods for manufacture of a medicament intended for the treatment and/or prevention of an NFB-regulated inflammatory response. In addition, methods for determining whether a test compound inhibits or enhances the activity of NFIF polypeptides are provided.

15